

## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



8  
84  
5

# AGRICULTURAL Research

June 1958

6/12



CAN PLANT NUTRITION AFFECT  
INSECT CONTROL?

*Page 4*

HOW COTTON SOILS

*Page 8*

FINDING FUNGI THAT HELP TREES

*Page 3*

GAINS THROUGH ROP

*Page 10*

U. S. DEPARTMENT OF AGRICULTURE



# AGRICULTURAL Research

Vol. 6—June 1958—No. 12  
(May 1958 issue erroneously designated Vol. 7)

## CONTENTS

### CROPS AND SOILS

- Finding Fungi That Help Trees..... 3  
Can Plant Nutrition Affect Insect Control?.. 4

### FOOD AND HOME

- Will a T-Shirt Keep Its Shape?..... 5  
Tougher Coating From Tung Oil..... 6  
Role of Fat in Nutrition..... 7  
How Cotton Soils..... 8

### LIVESTOCK

- Look for Gains Through ROP..... 10  
Saving Ranch Minks..... 12

### DAIRY

- We're Producing More Silage..... 12  
A Cow's Use of Energy..... 13

### FRUITS AND VEGETABLES

- New Machine Harvests Prunes Efficiently... 14

### AGRISEARCH NOTES

- Drying Rice Efficiently..... 15  
Pasture Plus Stilbestrol..... 15  
Control of a Potato Pest..... 15  
Pioneering in Nutrition..... 15  
New High-Sugar Sorgo..... 16  
Chelates Nourish Fruits..... 16  
Bagasse Shipping Boxes..... 16

Editor: J. F. Silbaugh. Managing Editor: J. R. Deatherage. Contributors to this issue: M. S. Peter, D. R. Klein, E. Evers, V. Bourdette, G. F. Snell.

Information in this periodical is public property and may be reprinted without permission. Mention of the source will be appreciated but it is not required.

## Future

We're all familiar with the achievements of United States agriculture over the last few decades—how a farm population decreasing by nearly a third has increased farm output by two-fifths since 1940, how net farm income for the country as a whole has more than doubled since the 1937-41 years. Farmers have enlarged their farms, further mechanized their operations, improved their practices, and adopted modern technology.

Yes, the past is remarkable. What about the future?

The long-range opportunities for farming look favorable.

Readjustments will be necessary, of course. It looks as though we may be struggling with overabundance for a decade ahead. We were producing in 1957 at an annual rate about equal to what we may be needing 4 or 5 years from now.

Looking further ahead, however, we see the possibility of needing 35 to 45 percent more farm output (above 1957 levels) to meet projected demand in 1975. It's estimated that we'll have 230 million people to feed in this country by that time. And urban development will bid for farmland and labor.

Indications are that we will be able to meet the expanding need for food and fiber. But to do that, we must look for still greater improvements in production efficiency. Present research suggests that we are going to find these improvements. This will reduce a farmer's risk, improve his opportunities, and make him less vulnerable on price declines.

The most important possibilities seem to lie in raising the output per acre and animal. Learning how to utilize our foods better and preventing crop losses before and after harvest offer still other worthwhile possibilities.

Economists tell us farms will become larger and more efficient. Farmers will buy more and more of their production factors and become more specialized. Cash outlays will increase.

We have some backlog of known research results that can be applied to increase output whenever it proves profitable. But continued increases in production efficiency depend largely on a vigorous, imaginative research effort involving both basic and applied investigations. Only with such support will we be able to supply added production as the needs arise.

*Agricultural Research* is published monthly by the Agricultural Research Service, United States Department of Agriculture, Washington 25, D. C. The printing of this publication has been approved by the Bureau of the Budget, September 16, 1955. Yearly subscription rate is \$1 in the United States and countries of the Postal Union, \$1.35 in other countries. Single copies are 15 cents each. Subscription orders should be sent to the Superintendent of Documents, Government Printing Office, Washington 25, D. C.

AGRICULTURAL RESEARCH SERVICE  
United States Department of Agriculture






MYCORRHIZAE formations on slash-pine seedlings are examined by plant pathologist J. G. Palmer, working in greenhouse at Agricultural Research Center, Beltsville, Md.



ROOTS of the pine, removed from the soil, reveal fungi covering entire sections. Formations help seedling to grow much more rapidly.



THREADLIKE filaments of ectotrophic fungus, used for experiments, penetrate between the cells, and radiate into a large volume of the soil. Endotrophic fungi, on the other hand, penetrate inside the cells of the seedling roots.

## FINDING FUNGI THAT HELP TREES

*We're looking for the most beneficial organisms and for ways to inoculate the soil*

**MANY ORGANISMS** hinder plant growth, but some formations, such as mycorrhizae, directly benefit growth.

Mycorrhizae are formations of fungi and roots. The fungi penetrate the roots of trees and shrubs and help trees absorb the nutrients from the soil. Trees will grow, of course, if climatic and soil conditions permit, but growth will be better in the more infertile soils when natural development of mycorrhizae is best.

USDA plant physiologist Edward Hacskeylo and plant pathologist J. G. Palmer want to find which mycorrhizal fungi are most beneficial to tree growth. There are two kinds:

1. Ectotrophic fungi (AGR. RES., March 1956, p. 6) have threadlike filaments that form covers around the roots, penetrate *between* the cells, and radiate into a large volume of the soil. This enables the trees to obtain an

increased amount of the nutrients that are available in the root zone.

2. Endotrophic fungi have filaments that penetrate *inside* root cells. The roots digest the fungus and absorb it as additional nutrients.

The ARS researchers set out to establish tree seedlings and inoculate them with different kinds of fungi to determine which are the most efficient. But first, it was necessary to find a method of removing all mycorrhizal fungi from the soil so that controlled pure culture inoculations would be insured for the experiments.

### Treatment stops formation

Studying the effects of a variety of chemicals on ectotrophic mycorrhizae, scientists recently learned that pretreatment of the soil with methyl bromide at 1 pound per 25 square feet prevents formation of mycorrhizae

on Virginia and slash pines during the first growing season.


### Chemical increases growth

Methyl bromide also resulted in increased growth of seedlings. They were planted after fumigation, since the chemical is poisonous to living plants. Reduction of other organisms in the soil probably reduced competition for nutrients, enabling the seedlings to grow faster.

Increased growth of the plots treated with methyl bromide continued during the second growing season, but mycorrhizae appeared on the seedlings. Possibly residual filaments or spores in the soil, or airborne spores of mycorrhizal fungi, or both served as inoculum for the plots.

Since soil becomes repopulated naturally by uncontrolled mycorrhizal fungi by the second season,





**SPORES** of *Amanita muscaria* fungi come from mushrooms growing near some of trees. These spores will be scraped from the dish and used for inoculation experiments.

future studies with specific forms will be undertaken through controlled inoculations during the first season. At this time, other fungi are absent, or present in only small numbers.

#### Inoculation method sought

Researchers are now looking for the best method of inoculating soil. They are experimenting in the laboratory with organisms on grains. If organisms grow successfully on the

grains, they will be sown. Still another method of inoculation under study is use of fungus spores from mushrooms growing around the trees.

If inoculations are successful, and the best organisms can be traced, then the problem of transplanting them to areas where they may become established to aid trees will have to be solved. The effect of different mycorrhizal fungi on other species of trees will also be given careful study. ☆

## CAN PLANT NUTRITION AFFECT INSECT CONTROL?

**Pests seem harder or easier to kill,  
depending on fertilizer diet of the host crop**

■ WHEN PLANTS are supplied with amounts of nitrogen, phosphorus, or potassium providing good growth, spider mites feeding on these plants are easier to kill with malathion.

Mites also are easier to kill when feeding on plants that have received a low level of phosphorus.

But mites are more difficult to kill when too much or not enough nitrogen or potassium nutrient is applied.

This is the result of preliminary experiments conducted at USDA's Agricultural Research Center, Beltsville, Md., with two-spotted spider mites on pole-lima beans. This may mean insect and mite control could be easier or more difficult depending upon the nutrition of the plants.

#### Uses of findings suggested

Possible applications of this information might be made in regulating the kind and amount of fertilizer to use for maximum effect on insects and mites, as well as for a maximum crop. Difference in susceptibility under various nutrient levels may possibly explain reported resistance to insecticides in insects and mites. Perhaps

control practices could be timed to take advantage of the seasonal variations in the level of nutrients in the tissues of the plant.

ARS entomologist Thomas Henneberry is in charge of the experiments, being conducted in cooperation with plant physiologist N. W. Stuart.

#### Workers study interactions

Nutrient solutions are applied to the bean plants the first day after potting, and twice a week thereafter. Different levels of nitrogen, phosphorus, and potassium are being tried independently and together.

With high levels of phosphorus and all levels of nitrogen, irrespective of the potassium levels, the mites are harder to kill. With high levels of potassium and high levels of nitrogen, irrespective of the phosphorus levels, the mites are likewise harder to kill. These and other interactions require further study for full understanding.

Researchers are also carrying out investigations in which mite susceptibility will be measured in relation to plant-growth variation produced by temperature and light treatment. ☆



**NUMBER** of mites on leaves from each plant is tabulated by entomologist Henneberry to determine different survival counts of mites on pole-lima beans fed different diets.




**MITES** are easier to kill with malathion on healthy plants (left) fed enough nitrogen for good growth. But mites are harder to kill on plants (right) supplied no nitrogen.



## Will a T-shirt Keep Its Shape?



**ROUND LOOPS** before washing (inset) show relaxed fabric. Shirt retains shape after 20 launderings, tumbler dryings. (Ink marks enclose area originally 10 inches square.)



**ELONGATED LOOPS** before washing (inset) indicate shirt will widen, shorten (note ink marks). Sleeves and neckband, cut crosswise on fabric, ruffled from excessive stretching.

**The shape of the loops, or stitches, tells whether the garment will shrink or stretch**

**■ CAN A PURCHASER TELL** whether T-shirts or other plain knit cotton garments will keep shape after laundering?

Research in the USDA Institute of Home Economics provides a key. It lies in the geometry of the knit. That is, a garment will have stable dimensions if the fabric loops (stitches) are round or slightly wider than long and there are more courses (loops counted lengthwise) per inch than wales (loops counted crosswise) per inch.

Stitches are pulled lengthwise during knitting. Round loops in a garment indicate these stitches have already been relaxed to natural shape and there will be little further change in dimension. If the loops are still elongated, however, they will relax during laundering, and the fabric will become wider and shorter.

These findings confirm earlier USDA studies (AGR. RES., November 1953, p. 13; May 1957, p. 11). These showed that stretching and shrinking during laundering were caused largely by changes in loop shape, whether the fabric was tricot or plain knit, or whether it was made of cotton, rayon, acetate, or nylon fiber.

ARS textile physicists S. H. Roberts and H. M. Fletcher included shirts of 33 different brands in the recent study. These included many sizes of yarn and loops of various width-to-length ratios. The researchers marked the shirts with three 10-inch distances in each direction and counted the wales and courses per inch between these markings. After each washing the distances between markings were measured; after five washings the wales and courses were counted. Differences in measurements and wale and course counts between washed and unwashed shirts showed the effect laundering had on the garments.

Shirts with loops having a width-to-length ratio of less than 1.0 before laundering shrank a great deal in length when washed and many stretched excessively in width. Garments with a ratio greater than 1.0 before laundering retained their shape and shrank only about 6 percent in both length and width when dried flat on a screen. After laundering, the width-to-length ratio of loops in all the garments approximated 1.2. Drying in a tumbler drier increased the amount of shrinkage in the garment. ☆



# *Tougher Coating from* **TUNG OIL**

**Greater serviceability and resistance to fungus damage have been built into new varnish**

■ RESULTS OF RECENT research in tung oil show great promise in paints and varnishes with hoped-for properties such as greater toughness and resistance to water, alkali and mildew.

Tung oil, because of its rapid-drying properties, has long been of interest to the paint and varnish industry. But the highly reactive character of the oil requires different techniques

than those used with soybean, linseed, and other drying oils.

Important exploratory work has been done during the past 3 years by L. L. Hopper, Jr., fellow of the Tung Research and Development League, and chemist L. A. Goldblatt at USDA's Southern Utilization Research and Development Division, New Orleans.


Many modern protective coatings are made with alkyd resins, or synthetic oils, usually a combination of a polyfunctional alcohol and acid. Recently, an alkyd in which all the drying oil is tung oil, was prepared on a

pilot-plant scale for a drum-coating enamel. It is now being evaluated by a major manufacturer.

Epoxy resins seemed to hold even more promise than the alkyds. These are synthetic resins obtained by the condensation of phenol, acetone, and epichlorohydrin. However, technical literature indicated it was difficult or even impossible to prepare epoxy resin esters with tung oil.

This served as a challenge, for formulations with epoxy resins offer extremely good adhesions, toughness, chemical resistance, and production

**BAKED ENAMEL** made with tung oil shows its toughness by bending sharply without cracking (right), whereas standard enamel on other metal strip cracked under stress.



**FUNGUS CULTURES** are prepared for test on varnishes. Some wooden blocks were coated with ordinary varnish, others with experimental varnish made with tung oil and fungicide. Microbiologist Ruth Y. Mayne inoculated specimens with fungi and will incubate them to see whether fungi can grow on the varnishes.




**TEST BLOCKS** have been kept for 7 to 10 days in a warm, moist chamber under conditions that frequently cause varnishes to become moldy. Inspection will show whether experimental varnish can withstand fungi.




of a hard, glossy film. Satisfactory results were finally obtained through the use of zinc resinate as a catalyst for the ester exchange. The zinc resinate also served to improve the water and alkali resistance of the finished coating. This is of great significance for many uses—for example, as finishes for washing machines.

Still another development is the inclusion of long-term, built-in fungicidal properties. Tung oil actually has excessive drying power while various fungicidal compounds that might be used are essentially nondrying. The scientists successfully combined these two characteristics and developed a potential fungicide coating which is now being evaluated. Extensive evaluation work on this and other new coatings is being planned in cooperation with industry. ☆

**GRAPHIC EVIDENCE** of effectiveness of fungicide in tung-oil varnish is shown here. Top square was coated with ordinary varnish, bottom square with fungicidal varnish.



## The Role of



## in Nutrition

■ THE COMPLICATED CONTRIBUTION of fat in nutrition is just beginning to be recognized and many researchers are working on various aspects. The USDA Institute of Home Economics is among those conducting research to find out the amounts and kinds of fats required for optimum health. ARS researcher Ruth M. Leverton sums up our present knowledge about the need for fat and the part played by linoleic acid as follows:

Fats are an important kind of food for all of us. They add variety and flavor to many foods, are concentrated sources of energy, carry vitamins A and D, and supply fatty acids that are essential for growth and health. Also, fats spare protein so that it is available for its specialized functions. Fat tissue in the body is important for support, protection, and insulation of vital organs and areas. In other words, fat is a *normal constituent* of food. And the body's use of fat for fuel is a *normal process*.

Linoleic acid is of particular importance. It is an essential unsaturated fatty acid and part of many fat molecules. It is needed for such vital functions as keeping the skin healthy.

Because the body cannot manufacture it, linoleic acid must be supplied in foods we eat. Common foods that contain appreciable amounts are the natural oils from corn, cottonseed, and soybean. Peanut oil and poultry fat have less linoleic acid, olive oil and pork fat still less. The fats of beef, lamb, milk, veal, and cocoanut oil contain very little linoleic acid. Margarine and shortenings differ widely in linoleic-acid content, depending on raw materials used and extent to which they are hydrogenated.

There is some evidence that fats with a high content of linoleic acid may help to lower blood cholesterol levels. Cholesterol is synthesized in the body and is a normal constituent of blood. There have been some indications that a high level of cholesterol is associated with atherosclerosis—a type of arteriosclerosis in which fatty material is deposited along the inner lining of artery walls. However, we lack evidence at this time that lowering blood cholesterol reduces the occurrence of atherosclerosis.

To most of the questions being asked on fat requirements and metabolism, the effect of different processing procedures on the nutritive value of fats, and the relation of fat to heart disease, we must answer that we do not know. But much research is in progress, and the practical results of these investigations will be applied as soon as possible to improve the Nation's food and its health. ☆



# HOW COTTON SOILS

**Findings are helping scientists move ahead in search for effective soil-resistant finishes**

■ WE HAVEN'T YET come up with a cotton that *completely* repels soil, but we have come a long way in finding out exactly how and why cotton *does* soil and what can be done about it.

Present work at USDA's Southern Utilization Research and Development Division, New Orleans, is aimed at improving cotton's qualities, with much effort going into improvement of resistance to dry soiling.

It's necessary to understand the mechanism of dry soiling before really effective soil-resistant finishes can be developed. Development of such hard-to-soil cottons would mean longer life and lower upkeep costs, especially for hard-to-laundry drapes and upholstery fabrics.

Gray and chemically modified cottons were artificially soiled in con-

trolled tests with colloidal and activated carbon, floor soil, and a synthetic soil and clay. Soiled cottons were examined by oblique lighting and in an electron microscope.

## **Fabric spaces hold the soil**

The geometry of the fabric—the way the fibers are physically put together—plays a major role in determining the distribution of the soil. The soil particles are mechanically and unevenly trapped between the fibers near the surface of the fabric and on the fiber surface because of the natural irregularities, grooves, and ridges of many cottons.

ARS chemists V. W. Tripp and Anna T. Moore, optic physicist Blanche R. Porter, and physical scientist Mary L. Rollins found a wide

range in the pickup of soil by cottons treated with typical finishing agents.

The hard, fine "particulate" finishes, for instance, greatly improved the soil resistance of the cotton. These granular finishes—which impart a colorless, sandpaper-like coating to the fibers—probably act by filling in the potential sites of soil attachment so soil particles can't get in. Examples of these finishes are colloidal silica and alumina, applied to the surface of the fabric from water suspensions. Another finish—perfluorodecanoic acid—gave significant resistance to soiling without changing the fiber surface.

Fabrics that were treated with a soft resin (like acrylic polymer latex emulsion) soiled easily and quickly. Soil particles got into the soft coating



OBLIQUE LIGHTING clearly reveals the pattern of soil entrapment between fibers of soiled, untreated bleached fabric. Low magnification was used with this method.





easily and embedded themselves firmly, a fact perhaps noted by persons who wear resin-treated fabrics.

The extent of the soiling apparently depends upon how soil gets into a fabric and how it is held there.

### Fiber folds also trap dirt

The natural irregularities of the fiber surface—folds or wrinkles, grooves, and ridges—serve to trap and imprison the dirt. These irregularities persist through routine chemical modifications such as mercerization—treatment with caustic alkali to make cottons stronger, silkier, easier to dye. Irregularities also persist through newer and more specialized modifications such as partial acetylation and cyanoethylation, resin impregnation, and some types of additive finishing (AGR. RES., October 1957, p. 15; June 1957, p. 15; October 1955, p. 16). And the more dirt in a spot, the more will collect on top of it. Size of soil particles helps determine the extent of soiling, too. Finer particles

get into the fiber surface deeper than large particles ordinarily do.

Researchers found that dirt also attaches to smooth, regular fiber surfaces. This shows that soil adsorption is due to chemical attraction as well as mechanical entrapment.

### Chemical treatment best hope

Modifications of the fiber surface through scouring (which gets rid of cotton's noncellulosic coating) or by carboxymethylation (which roughens

the surface) have little effect on the distribution of soil on the fiber.

The best present antisoiling agents are additive finishes or chemical modifications that tend to lower the energy of the fiber surface. Perhaps really effective soil resistance can be built into future cotton by combinations of various treatments. Much work remains to be done, however, on surface-energy effects in soiling of cotton, and on adaption of modifications for commercial use. ☆


**ELECTRON** microscope is utilized by chemist Anna T. Moore to examine location of soiling agents in fibers from chemically treated and untreated fabrics. Microscope magnifies greatly, hence greater detail is visible. Circular inset shows unsoiled and soiled fibers at 500 X magnification.




## UNSOILED




## SOILED



Fibers from a fabric not bleached or otherwise chemically modified



Fibers from a fabric modified by soft acrylic polymer latex emulsion



Fibers from a fabric modified by treatment with commercial formulation

**(10,000 X MAGNIFICATION)**



# Look for Gains Through **ROP**

## Virginia's research-based improvement effort pays off in more efficient beef production

■ VIRGINIA'S BEEF CATTLE have earned a solid reputation for predictability and excellence, thanks to USDA-State research and marketing procedures.

Current improvement in the State's cattle is being made through a statewide record-of-performance (ROP) testing program by the recently organized Virginia Beef Cattle Improvement Association. Its work is based almost entirely on research results from several years of beef testing at the Beef Cattle Research Station, Front Royal, Va. (AGR. RES., April 1955, p. 3).

Researchers there have found that a bull's early performance record is useful in predicting similar performance (rate of gain, conformation to good carcass type, and feed-use efficiency) in his offspring. No longer need a breeder wait until a bull is 2 or 3 years old before finding out his potential value as a future herd sire.

### Good showing spurred breeders to organize

Limited testing in Virginia and elsewhere confirmed the practicability of judging a bull's breeding potential by calfhood tests. Virginia Agricultural Experiment Station researchers, the Virginia Division of Markets, and breeders realized the importance for herd improvement of recognizing outstanding bulls early. So in 1955 they organized State breeders for ROP testing of beef cattle. This was

the first such statewide plan. Similar groups have since been started in a number of other States.

What is performance testing and how does it operate?


It's the evaluation of inherited differences in ability to grow rapidly, to mature early, to use feed efficiently, and to develop the desired type of carcass. It's done by measuring gain before and after weaning, feed requirements per pound of gain, and grade (or type desirability). Confidential data are provided to association members on *all* progeny of each bull and cow in member herds.

### Professional and record services are given

When a breeder joins the association—which he can do if he has 25 or more purebred or grade cows—he gets a code number for his herd and an IBM card for each cow. When a cow calves, the breeder gives the calf a number to identify it, records the number, the calf's sex, birth date, and weight, and the identification number of its sire and dam. He mails a card on each calf to the State supervisor in Blacksburg. Just before weaning time a team visits the farm to weigh and grade his cattle.

Breeders get from the State office the performance record of each calf, showing age, weight, type score, average daily gain, and index value. All calves by one sire are grouped and their average performance shown. At the end of the year, each breeder gets a performance record of the progeny of all organization sires.

Thus, a breeder can better evaluate the worth of each animal in his herd. He can compare gains and type scores



CALF in upper left picture will soon tell through performance tests not only his own intrinsic worth as a herd sire but also, in part, the value of his dam as a breeder. Value of such testing is shown by bulls in corner picture. They may appear similar, but weanling calves of bull at left were worth \$12 per head more than other bull's calves. Weighing and grading—important part of ROP testing—are done on farm (below) when animals are 4 to 10 months old. Virginia association frequently provides portable scales.







**BRISK BIDDING** by livestockmen and fast sale of bulls at recent Front Royal auction is indication of value of ROP-tested cattle. Bulls were bought with confidence by beef breeders anxious to improve herds. Animals brought good prices.

of calves sired by different bulls. He can compare the performance of all cows in his herd. He can see how his herd compares with other association herds in his State.

### **Records point out best animals in herd**

This information enables a breeder to decide which bull is doing the best job of improvement, which cows are producing the best calves. It provides a guide for culling the herd and for deciding which heifers to save for herd replacements. Such detailed information plus the breed-

er's personal knowledge of his cattle are the best basis for improving growth rate and conformation.

There are about 150 herds in the Virginia Beef Cattle Improvement Association. There's a membership fee and a small weighing and grading charge. But these are more than offset by greater returns from the cattle.

Researchers, as well as herd owners, believe performance testing of beef cattle is here to stay—improving efficiency of beef production, and giving the consumer a better meat product at a somewhat lower cost. ☆



**PREMIUMS** of \$25 per head were paid by Dunbar Oswald (right in above picture) for 100 ROP cattle to stock a South Carolina farm. Beside him, Thomas Marlowe, executive secretary of Virginia Beef Cattle Improvement Association, negotiates the sale with seller Prescott Carter. At right, association-tested cattle at recent Front Royal auction are discussed by State-Federal researchers (l. to r.): Kenneth Bovard, Marlowe, Charles Kincaid, George Litton, and Everett Warwick.





# *Saving Ranch* **MINKS**



■ WESTERN RANCH-RAISED MINKS face the threat of extinction from virus enteritis if the disease continues to run its present course, say USDA researchers. Virus enteritis is a deadly intestinal disease that usually kills affected animals. There has been a recent sharp rise in outbreaks of the disease in Wisconsin, Michigan, New York, Illinois, and Massachusetts.

There's no way to treat infected animals. But growers and veterinarians can take measures to prevent or check the disease, according to ARS veterinarians D. Burger, R. K. Farrell, and J. R. Gorham, who are stationed at the Washington Agricultural Experiment Station, Pullman.

Two types of vaccine are available. Perhaps the most effective is one prepared from spleens of animals dead of the disease. The other is a commercial vaccine used against feline panleucopenia—blood disease of cats. Al-

**CAGED, ranch-raised mink are especially susceptible to new disease. It spreads easily from pen to pen and kills the kits rapidly. Adults survive.**

though this vaccine isn't specific for virus enteritis, it does give beneficial results. First dose of preventive vaccines should be given about the last week of June followed by a second dose administered 10 days later.

Virus enteritis in minks occurs mostly in July, August, and September, and even up to pelting time. June outbreaks, fortunately uncommon, are quite difficult to control. Kits are most susceptible; adults are fairly resistant and seldom die.

The disease may reappear each summer on one ranch. Losses of half the kits are common. Exact mode of

spread is unknown. Carrier minks and maybe cats may bring the disease to a clean ranch. Birds and insects may carry the disease, too. The disease pattern is similar to distemper. It spreads easily to nearby pens, but new cases may be noticed many yards away from known infected pens.

First sign of the disease is refusal of infected animals to eat. At autopsy, these animals show marked inflammation of the intestines. The lining is swollen and cells are deteriorated. The spleen is enlarged and dark and may show signs of hemorrhaging. Lungs and liver look normal. ☆

## **DAIRY · DAIRY · DAIRY · DAIRY · DAIRY · D**

### **WE'RE PRODUCING MORE SILAGE**

■ SILAGE PRODUCTION KEEPS GROWING, and harvesting methods keep improving. Horsedrawn cutting sleds were used when silage was first reported, on a small scale, in this country in 1875. But field choppers have harvested over 80 percent of the silage in recent years.

A USDA survey indicates that about one-eighth of our farms used silage in 1955. The study, conducted by ARS and Agricultural Marketing Service on 29,000 farms, shows an increase in silage making and in number of farms using silage. This trend is pronounced everywhere except in the Lake and Northeastern States.

Production in 1955 ran 73 million tons—one-third more than in 1951. The 600,000 farms making silage in 1955 represent a 20-percent increase in 5 years.

The study, which includes corn, grass, sorghum, and other silage, shows greatest use where dairying is important. Corn accounted for 74 percent of all silage.

Tower silos stored 70 percent of the corn silage, with the balance in trench, bunker, and temporary silos.

Grass silage, totaling 9.3 million tons for the United States, was made from hay crops, small grains and pasture clippings. The main crops used were alfalfa, clover, soybean, cowpeas, vetch, and sudan grass.

About one-third of the grass silage was treated with a preservative. Sodium metabisulfite was used as the only preservative on about 45 percent of these farms. Molasses preservative was used alone on 23 percent.

Sorghum silage totaled 9.4 million tons, with the bulk in Kansas, Texas, Missouri, and Oklahoma. Drought-resistant qualities make sorghum a leading silage crop in much of the southern and central plains area.

"Other silage" (byproducts of canning and processing crops, and other products) totaled 1,160,000 tons and was most important in Northeastern and Pacific States.

Mechanical unloaders were used for removing only 4 percent of all kinds of silage that was stored in tower silos in 1955. Six percent of the trench and temporary silos were constructed to allow self-feeding. ☆



*Regardless of this cow's activity on pasture the gases she exhales are measured and sampled by a device that's helping us learn more about*

## A COW'S USE OF ENERGY

■ A PORTABLE METER has been strapped to a cow's back to measure her exhalations. USDA scientists are using the information to find how much energy a cow uses on pasture.

D. R. Waldo, ARS dairy husbandman, is conducting the experiment at the Agricultural Research Center, Beltsville, Md. He wants to find out why a cow uses more energy to maintain her body in a pasture than in a barn. And he wants to find out the extent of this energy expenditure.

Two test cows roam the pastures in a natural environment. A tracheal

transection has been performed so that air going in and out of each cow's lungs will pass through a plastic valve and hose. The volume of the respiratory gases the animal exhales is measured in liters on the meter. And a percentage of the gas is collected continuously in a bag harnessed to the cow. The urine and feces are also being collected in a container, which is harnessed to each of the cows while she is on pasture.


Gas samples will be analyzed to determine amount of oxygen, carbon dioxide, methane, and nitrogen.

When the amount of gases animals burn and the amount of nitrogen in the urine are known, the heat production (amount of work animals do) can be computed in terms of calories.

Recordings will be made under different environmental conditions—varying in temperature, wind velocity, and sunlight. The amount of exercise the animals take will change according to the kind of pasture. If the grasses are short, the cows will obviously have to walk greater distances and take more bites than if the grasses are high. The level of food intake by the cow will be estimated to determine how heat production will vary under different conditions.

Both this pasture study to determine the amount of energy a cow uses and a study to determine the nutritional value of feed (AGR. RES., May 1958, p. 8) are part of the energy-metabolism laboratory activities. ☆

AMOUNT of gas cow exhales is indicated on dial resembling meter on gasoline pump. Bag on other side collects gas samples. Containers gather urine, feces. Two cows will roam pasture during tests. Checking equipment are D. R. Waldo (left), and W. P. Flatt, head of the energy-metabolism laboratory.





# NEW MACHINE HARVESTS PRUNES EFFICIENTLY

*Device eliminates stoop labor  
in an important crop*

■ THE HIGH LABOR COST may have been taken out of harvesting prunes for drying, through cooperative research by USDA and California Agricultural Experiment Station.

Harvesting has been a major problem to prune growers, particularly in Washington, California, Oregon, and Idaho, where more than 400,000 tons of prunes are produced and dried annually. In usual practice, prunes are not picked directly from the trees. Harvesting begins as the fruit ripens and falls to the ground. Then crews are hired to pick up the prunes by hand. Occasionally, the trees are shaken to dislodge ripe prunes that have not fallen.

In recent trials at Davis, Calif., agricultural engineers P. A. Adrian, of ARS, and R. B. Fridley and A. A. McKillop, of the State station, have successfully tested a pilot mechanical harvester. This 1-man, self-propelled machine harvested a 20-inch swath and picked up about 1,000 pounds of prunes per hour. That replaces about four hand laborers ordinarily used.



As a result of these tests, USDA will apply for a public patent on the principle used in the machine. However, research is continuing on this and other models in an effort to develop an even more efficient harvester. The engineers hope to interest manufacturers who will produce it and place it in the hands of the growers.

The pickup device on the experimental harvester resembles the wringer of a washing machine. Two small rollers, heavily padded with



**HARVESTING MACHINE** by researchers of USDA and California Agricultural Experiment Station picks up fallen prunes under trees. This displaces the costly and thankless task of picking them up by hand.

**COUNTER SPIN** enables the soft rolls to gently grip prunes on the ground, raise them into machine.



**SOFT AUGURS** in front sweep fruits toward center of swath. Pickup rolls are at height to reach but not crush fruits.

sponge rubber, rotate to pass objects between them as the machine moves over the ground. The fallen fruit is pulled into the space between the rollers and tossed back onto a conveyor belt that carries the prunes to a box at the rear of the machine. For most efficient operation, the orchard floor must be level and free of stones, broken branches, and other debris.

It was found that at least two runs through the orchard are necessary to pick up all the prunes in growing areas where not all the fruit on a tree matures at the same time.

For areas where a tree's entire crop does ripen at once, USDA agricultural engineers are experimenting with the use of mechanical tree shakers for removing all of the fruit. ☆



## Drying rice efficiently

Raising temperature of a rice dryer and putting the rice through it more times but at a faster rate has boosted capacity of conventional equipment up to 58 percent, head yield by 2.5 percent, and total rice yield by 1.5 percent in USDA-grower studies. Drying costs were reduced, too.

A drying temperature of about 137° F., up to 6 passes through the dryer, and a drying rate of about 1,760 bags per hour proved to be about optimum in the plant-scale tests by the ARS Western Utilization Research and Development Division, Albany, Calif., and the Sutter Basin Rice Growers Cooperative at Knight's Landing, Calif. This method removed less moisture per passage through the dryer and reduced damage to the rice.

## Pasture plus stilbestrol

Stilbestrol—hormone-like chemical now received by three-quarters of the beef cattle in the country's feedlots—may also help improve summer gains of yearling steers on native range, State-USDA research shows.

Yearling Hereford steers were treated with 30 milligrams of the chemical, implanted under the skin of the ear, in grazing experiments at the USDA Northern Great Plains Field Station, Mandan, N. Dak.

For the 140-day feeding test on moderately grazed pastures, implanted



steers averaged 48 pounds more per head than untreated steers. Not all of the steers in this test reacted the same to stilbestrol. Some treated animals

gained less than some of the controls. But the variation wasn't great enough to offset the extra gain, averaging nearly 21 percent, made by the steers implanted with stilbestrol.

In a similar test—but on heavily grazed pastures—implanted steers kept on gaining even when some of the control animals lost weight because they couldn't get enough feed.

Although implanted steers gained much more than untreated steers on both moderately and heavily grazed pastures, the gain advantage due to having a *d e q u a t e* feed was much greater than that due to stilbestrol.

We need to learn what effect can be expected from a second stilbestrol treatment after steers go to the feedlot for finishing. Some research suggests that stilbestrol produces greater gains only once. Many Corn Belt farmers are thus reluctant to buy feeder steers that were implanted while on pasture, fearing that further stilbestrol during drylot feeding may be ineffective. Selling price of implanted range steers might therefore be reduced enough to take the profit out of using stilbestrol to increase gains.

## Control of a potato pest

The green peach aphid that attacks potatoes in the Pacific Northwest has been controlled under experimental conditions with Thiodan. Two years of USDA testing with various formulations and dosages showed Thiodan gave excellent control (90 to 99 percent) when applied by aircraft. The insecticide was found superior to demeton, Bayer 19639, trithion, thimet, malathion, guthion, American Cyanamid 4124, and parathion.

ARS entomologists working at Union Gap, Wash., used an emulsion spray made of 1 pound of Thiodan in

10 gallons of water per acre. This was superior to the same dosage prepared in kerosene, or formulated as a 60-mesh granule dust, or as a 325-mesh impregnated dust.

The insecticide killed relatively few of the larvae or adults of the *coccinella*, a beneficial insect that feeds on the aphids. Thiodan also controlled the tuber flea beetle and the important Colorado potato beetle.

## Pioneering in nutrition

New insight into nutrition of living cells—how nutrients are used, requirements for various nutrients, and interrelationships between them—is the objective of the newly established USDA Pioneering Research Laboratory in Cellular Metabolism.

The researchers will use micro-organisms (bacteria, protozoa, algae, yeasts, and molds) as experimental tools for rapid exploratory work in nutrition. Radioisotopes will assist in following the course of nutrients as they are used by living cells. Some findings may be evaluated more specifically with animal cell cultures.

Micro-organisms are versatile tools in the investigation of a wide variety of food and nutrition problems. These living substances have so much in common with all other living organisms that learning how one species functions leads to new understanding of other species. Nutrients essential to higher animals are also essential to many micro-organisms. Many of the vitamins that function in the metabolism of micro-organisms also play essential roles in human and animal nutrition and, as far as is known, the individual vitamins perform similar enzymatic functions in all organisms.

This laboratory, part of the ARS Institute of Home Economics, will



**TES · AGRISEARCH NOTES · AGRISEA**

conduct its work at the Agricultural Research Center, Beltsville, Md. The laboratory's key scientists will include bacteriologists Howard Reynolds and Thomas P. O'Barr. These scientists have been carrying on microbiological research in the Institute of Home Economics for some time, including investigations of the thermal resistance of micro-organisms, microbiological assay procedures, and metabolic interrelationships among amino acids in micro-organisms.

### **New high-sugar sorgo**

Release of a new early maturing sorgo adapted for production under irrigation in California's Imperial Valley and in similar western areas is announced by USDA and California Agricultural Experiment Station.

Seed for Brawley—as the new sorgo is called—will be available from the Imperial Valley Field Station at El Centro for commercial production of registered seed in 1958. No seed will be made available by USDA.

Sorgo is commonly grown for use as a sirup and a livestock feed. Brawley, however, shapes up as a potential sugar crop for California, assuming we can develop a suitable processing



technique. Brawley ranks high in sucrose content and purity of juice and compares favorably with other

commercial sorgos such as Rex and Tracy in resistance to lodging.

Early maturity of the new variety makes possible two crops a year in the Imperial Valley. When so grown, Brawley was harvested earlier than Rex or Tracy, the height and total yield of the feed stalks was greater, and there was less lodging. Seed yield, however, averaged only 950 pounds per acre—about 100 pounds less than that of Rex and Tracy.

Brawley was one of the first varieties developed under a breeding program started in 1946 by USDA to produce new sorgos for culture under irrigation in the Imperial Valley. The new variety is a cross between selections from Rex, which has high lodging resistance, and from the white-seeded Collier, which has a high sucrose content, lodging resistance.

### **Chelates nourish fruits**

Chemicals known as chelates are being used in USDA-State research to deliver essential metals to plants under soil conditions that normally cause metal starvation.

A chelate is a soluble organic compound with a ring structure that holds a metal atom between two or more other atoms. This type of linkage strongly withholds the metal atom from various soil ions that would tie up soil metals in insoluble form in an open-end type of structure.

In Louisiana, pecan rosette, a zinc-deficiency disease, was overcome by soil treatment with zinc chelate.

Where irrigation has caused an alkaline condition in California and Texas soils, lime-induced chlorosis of



citrus trees was corrected by treating the soil with iron chelates.

Soil treatment in Washington with a zinc chelate corrected chlorosis of peaches and sweet cherries. In some orchard soils, arsenic has accumulated excessively in the soil from years of spraying. Chelated zinc has overcome arsenic toxicity there.

Chelates have been successfully used as foliar sprays, too. Pear chlorosis in Oregon was overcome by spraying with chelated manganese and iron. Chlorosis of apples elsewhere in the Pacific Northwest has yielded to a summer spray of zinc chelate used in connection with spring and fall sprays of zinc sulfate.

### **Bagasse shipping boxes**

Efficient can and bottle shipping boxes have been made from depithed sugarcane bagasse in USDA research.

Three-fourths of the bagasse boxes were equal or superior to comparable commercial boxes in laboratory compression tests, and half were equal or better in tumbler tests. Loaded bagasse boxes equaled commercial boxes in truck and rail hauls. Bagasse boxes have greater vertical stiffness but are more brittle.



74  
we

# AGRICULTURAL Research

## INDEX

OF VOLUMES 4,-5,-6

*July 1955 - June 1958*

U. S. DEPT. OF AGRICULTURE  
NATIONAL AGRICULTURAL LIBRARY

JAN 11 1966

C & R-ASF

U. S. DEPARTMENT OF AGRICULTURE



■ **AGRICULTURAL RESEARCH** reports on the problems and progress of research done by USDA scientists and cooperators to improve agricultural production, utilization, and marketing; on closely related regulatory activities that grow out of research; and on broad policies connected with such work. The magazine aims to provide accurate, readable information and interpretations of the latest and most significant developments to persons doing agricultural research or working with farmers.

Articles are listed by subject matter with cross references as necessary.



# A

AAventa nematocide. Feb. 1956, p. 12  
 Accidents, increase in farm fatalities. July 1957, p. 7  
 Acetoglycerides from fats, new uses. July 1955, p. 10  
 Aceto-oleins from fats, new uses. July 1955, p. 10  
 Acetostearins from fats, new uses. July 1955, p. 10  
 Additives to animal feeds. Oct. 1956, p. 10  
 Adjustments in farm operations, financing. Feb. 1957, p. 12  
 Advisory committees on USDA research:  
   Food and nutrition, recommendations. Jan. 1956, p. 16  
   New members, Research Policy. Dec. 1957, p. 16  
   Research and marketing. Nov. 1956, p. 15  
 Aerosol, improved can. Nov. 1957, p. 5  
*Agave lecheguilla*, source of cortisone. Mar. 1957, p. 15  
 Aircraft:  
   Apply pesticides. Sept. 1955, p. 14; Feb. 1957, p. 16;  
     Oct. 1957, p. 14  
   Fight forest fires. Nov. 1957, p. 7  
   Jet noise and livestock. Jan. 1957, p. 10  
 Alfalfa:  
   Creeping varieties. Oct. 1956, p. 4  
   Grown with grass in Northeast. Nov. 1955, p. 11  
   Moapa variety, aphid-resistant. Oct. 1957, p. 16  
   Silage for dairy cows. Dec. 1955, p. 16  
 Alfalfa aphid, spotted. July 1956, p. 16  
 Alkyl aryl sulfonate removes thrips from berries. Oct. 1956, p. 15  
 Aluminum foil for soil samples. Mar. 1957, p. 15  
 Aluminum and hydrangea color. July 1957, p. 15  
 Ames, Iowa, laboratory for animal disease—see USDA  
 Amino acids:  
   Amount in foods. Apr. 1958, p. 12  
   Human requirements for. July 1956, p. 11; May 1958, p. 6  
   Relation to liver fat and carbohydrate. Aug. 1956, p. 13  
 Anaplasmosis control in Hawaii. May 1957, p. 3  
 Ancestry determined by blood type. Sept. 1957, p. 8  
 Anesthetics used to measure animal's fat. Oct. 1956, p. 11  
 Aneuploids aid genetic studies of wheat. May 1956, p. 7  
 Antibiotics:  
   Chronic respiratory disease. Sept. 1957, p. 15  
   Coccidiosis control. Nov. 1956, p. 15  
   Control fungus diseases in vegetables. Mar. 1956, p. 10;  
     June 1956, p. 16  
   Feed additives. Oct. 1956, p. 10  
   International conference on. Jan. 1956, p. 15  
 Antisera for plant virus diseases. Feb. 1956, p. 3  
 Aphids:  
   Carriers of tristeza in citrus. Oct. 1955, p. 8  
   Spotted alfalfa aphid. July 1956, p. 16  
 Apples:  
   Color affected by light. Apr. 1957, p. 12  
   Handling made easier. Sept. 1955, p. 13  
   Insecticide ryania for pests. Dec. 1955, p. 16  
   Marketing in bulk boxes. Apr. 1957, p. 15  
   Mildew in East. Mar. 1956, p. 11  
   Pomace feed for cattle. May 1958, p. 3  
   Tetraploid sport for breeding big Winesap. Jan. 1956, p. 3  
   Trees strengthened with hay mulch. Jan. 1957, p. 14

Appraisals, tax, on soil earning power. Jan. 1957, p. 5  
 Apricots, new varieties. May 1958, p. 12  
 Ascorbic acid—see Vitamin C  
 AT, herbicide helps rebuild pastures. Nov. 1955, p. 11  
 Atomic energy:  
   Radioactive diets for poultry research. Oct. 1955, p. 13  
   Radioactive fertilizer for crop studies. Feb. 1956, p. 8  
   Sterile male technique for insects—see Nonchemical control  
 Atrophic rhinitis:  
   Cost of. Nov. 1956, p. 12  
   Raising disease-free hogs. Oct. 1955, p. 11  
   Rats may be carriers. Sept. 1957, p. 12  
 Attractant:  
   Flies deposit "fly factor." Aug. 1955, p. 15  
   For gypsy moth. May 1957, p. 8; June 1957, p. 7  
   Search for new. Sept. 1957, p. 7  
   Tomato, root knot nematode. June 1956, p. 12  
   See also Mediterranean fruit fly  
 Auctions, livestock, improving. Nov. 1956, p. 11  
 Avena test and cotton boll drop. May 1956, p. 6  
 Awards for researchers:  
   Bordon Award for Burmester. Dec. 1957, p. 16  
   Hoblitzelle awards to Olsen, Stephens, and Quinby. July 1955, p. 16  
   Presidential Award to Hendricks. April 1958, p. 3  
   Tom Newman Award for Burmester. Nov. 1956, p. 16  
   USDA awards. Aug. 1955, p. 3; Aug. 1956, p. 3; July 1957, p. 3  
 Awnings, canvas, treated to resist weather. May 1957, p. 15;  
   Mar. 1958, p. 16

# B

B vitamins—see Vitamin B<sub>12</sub>  
 Bacteria:  
   Culture collection. Apr. 1958, p. 8  
   Effect on silage quality. Aug. 1957, p. 10  
   Preserved with tetrazolium. Oct. 1956, p. 5  
 Bagasse, new uses for. July 1956, p. 4; June 1958, p. 16  
 Baking—see Bread, Flour quality  
 Bark:  
   Feeding of fruit trees. Sept. 1956, p. 11  
   Removing from logs. May 1958, p. 14  
 Barley, tests for stripe-mosaic virus. Feb. 1957, p. 8  
 Basting, pin or thread, in sewing. July 1957, p. 16  
 Bayer 19639, systemic for cotton insects. Feb. 1957, p. 10  
 Bayer 21/199, new systemic for cattle. Mar. 1958, p. 12  
 Beans:  
   Columbia Pinto, new variety. Aug. 1957, p. 15  
   Gamma-glutamyl-S-methyleysteine isolated. Jan. 1958, p. 15  
   Storage and eating quality. Dec. 1956, p. 12  
   Storage and yield. Dec. 1956, p. 12  
   Timing irrigation. Dec. 1956, p. 16  
   See also Lima beans  
 Beef:  
   Consumption increasing. July 1956, p. 15  
   Efficient use in quantity food service. Mar. 1956, p. 15  
   See also Cattle, beef  
 Bees, laborsaving equipment. Nov. 1957, p. 8



Beets—see Sugarbeets

Berries:

Wetting agent removes thrips. Oct. 1956, p. 15

See also Individual berry, Plant introduction

Berry boxes, fungicide prolongs life. Oct. 1956, p. 15

Biological control—see Nonchemical control

Biological warfare defense. Aug. 1957, p. 3; Sept. 1957, p. 13; Oct. 1957, p. 6

Blackberry introductions from South America. June 1957, p. 3

Blackhead parasite and cecal worm. Aug. 1957, p. 12

Blacklands (Texas), management. July 1955, p. 8

Blackspot on roses, controlling. Mar. 1956, p. 16; Mar. 1957, p. 16

Blackstrap—see Molasses.

Blight—see Fungi

Blind-seed disease of ryegrass, control. May 1956, p. 4

Bloat in livestock. Feb. 1957, p. 4

Blood types in cattle. Sept. 1957, p. 8

Blubarb jam from blueberries and rhubarb. Apr. 1957, p. 16

Blueberry:

Blubarb jam. Apr. 1957, p. 16

Breeding new varieties. Oct. 1957, p. 8

Canning, how to prevent clumping. Feb. 1958, p. 10

Boll weevils resist insecticides. May 1957, p. 7

Bollworm, pink—see Pink bollworm

Bread:

Freezing and storing. Dec. 1955, p. 13

From new varieties of wheat. Nov. 1955, p. 8

See also Flour quality

Breeding, general:

Plants. Aug. 1955, p. 2; Mar. 1957, p. 2; Apr. 1957, p. 2

See also individual plant or animal

Breeding resistant varieties:

Alfalfa. Oct. 1956, p. 4; Oct. 1957, p. 16

Beans. Aug. 1957, p. 15; Apr. 1958, p. 5

Berries. June 1957, p. 3; Oct. 1957, p. 8

Chestnuts. Jan. 1958, p. 15

Corn. Aug. 1955, p. 2; Nov. 1956, p. 3

Cotton. Nov. 1955, p. 5

Editorials. Aug. 1955, p. 2; Sept. 1955, p. 2; Apr. 1957, p. 2

Flax. Dec. 1955, p. 3; May 1958, p. 16

Grapes. Dec. 1955, p. 12; May 1958, p. 11

Grasses. Sept. 1955, p. 2; Oct. 1955, p. 6; Jan. 1956, p. 12; Feb. 1958, p. 6

Ladino clover. Apr. 1956, p. 7; Feb. 1958, p. 16

Lupine. Dec. 1957, p. 4

Oats. May 1956, p. 3; July 1956, p. 6; Feb. 1957, p. 16

Pines. Feb. 1957, p. 7

Potatoes. July 1955, p. 13; Aug. 1955, p. 16; Nov. 1955, p. 13

Rice. Dec. 1957, p. 15

Seed bank. Sept. 1956, p. 6

Sesame. June 1957, p. 4

Sorghum. Aug. 1956, p. 11

Soybeans. Feb. 1957, p. 11; Feb. 1958, p. 15

Sugarcane. July 1955, p. 7; Aug. 1956, p. 10; Nov. 1957, p. 10

Sweetpotatoes. June 1956, p. 11

Tobacco. Feb. 1956, p. 16

Breeding resistant varieties—Continued

Tomatoes. Aug. 1956, p. 12; May 1958, p. 16

Watermelons. Feb. 1958, p. 16

Wheat. Dec. 1955, p. 5; Feb. 1956, p. 7; May 1956, p. 7; Aug. 1956, p. 8

Brine shrimp for residue tests. Oct. 1955, p. 16

Broilers—see Chicken

Brooder—see Chicken

Brucellosis, cattle:

Age unrelated to susceptibility. Aug. 1957, p. 15

Eradication campaign evaluated. Nov. 1955, p. 4

Interstate movement regulations. Jan. 1957, p. 15

States certified. Aug. 1957, p. 16; Oct. 1957, p. 15; Nov. 1957, p. 15; Mar. 1958, p. 16; May 1958, p. 15

Brucellosis, hogs, false positive reaction. Feb. 1956, p. 4

Budget—see Food

Budworm in tobacco, controlled by DDT. Dec. 1956, p. 16

Buildings—see Housing

Bulbs, proper storage temperatures. Nov. 1957, p. 14

Bulk boxes for handling apples. Apr. 1957, p. 15

## C

Cake-freezing tips. Aug. 1957, p. 14

Calves—see Cattle

Camellia flower blight. Sept. 1957, p. 16

Candy:

Higher quality with longer life. Feb. 1956, p. 14

Use of fats. Nov. 1956, p. 16

Canning home:

Improved methods. June 1956, p. 15

Survey of practice. June 1956, p. 14

Canvas—see Cotton

Carbohydrates, relation to amino acids and liver fat in nutrition. Aug. 1956, p. 13

Careers in Government service. Apr. 1958, p. 2

Carotene:

From fungi. Nov. 1957, p. 16

In chestnuts. Apr. 1957, p. 16

Carpets—see Cotton fabric

Carrots, hydrocooling and packaging. May 1956, p. 11

Castorbeans, new varieties and harvesters. Mar. 1957, p. 12

Cattle:

Bloat, feedlot. Feb. 1957, p. 4

Diseases. Nov. 1955, p. 4; see also Anaplasmosis; Brucellosis, Emergency measures; Foot-and-mouth disease; Johne's disease; Lungworm; Laboratories, USDA

Dwarfism, X-rays can identify carriers. Jan. 1958, p. 4

Feed additives. Oct. 1956, p. 10

Feed, use of molasses to improve. Dec. 1956, p. 5

Fescue pasture and parasites. June 1957, p. 6

Fever tick. Dec. 1957, p. 8

Field-freeze-drying potatoes for feed. Oct. 1955, p. 10

Grub—see Grub, cattle

Heat tolerant. Sept. 1955, p. 11; Nov. 1955, p. 14; Dec. 1957, p. 10

Identification system. Dec. 1955, p. 15

Protection against heat, sun. Oct. 1957, p. 16

Rumen inoculation, calves don't need. June 1957, p. 11



## Cattle—Continued

Systemic for grubs helps control roundworms. Jan. 1958, p. 15

Systemic for grubs and screwworms. Mar. 1958, p. 12

See also Livestock, Stilbestrol

## Cattle, beef:

Apple pomace is nutritious feed. May 1958, p. 3

Breeding. Apr. 1956, p. 15; Feb. 1957, p. 15; June 1958, p. 10

Forest range in Florida. July 1957, p. 5

Interrupted gain. July 1956, p. 16

Investment in shelters. Aug. 1957, p. 7

Low-protein diet and calf crop. Apr. 1956, p. 13

Profitable feeding. Oct. 1955, p. 12

Protein needs of wintering calves. July 1957, p. 12

Stilbestrol—see Stilbestrol

Twin calves for research. May 1958, p. 15

Virginia Beef Cattle Improvement Assn. June 1958, p. 10

See also Range

## Cattle, dairy:

Blood types show ancestry. Sept. 1957, p. 8

Breeding. Feb. 1956, p. 13; June 1956, p. 4; Mar. 1957, p. 7; July 1957, p. 13

Butterfat record of Beltsville cow. Oct. 1957, p. 15

Cheaper ration for heifers. Apr. 1957, p. 16

Dairy feeds of the future. May 1956, p. 14

Dust cattle for hornfly. Apr. 1958, p. 15

Green-chop feeding can increase profit. Oct. 1957, p. 15

Heat tolerant—see Cattle

Herd behavior can reduce milk production. July 1955, p. 15

Home milk supply, produce or buy? June 1957, p. 10

Metabolism studies. May 1958, p. 8; June 1958, p. 13

New growth standards. Aug. 1955, p. 13

Pasture management. Mar. 1956, p. 14

Replacements, buying beats raising. Feb. 1957, p. 5

Stilbestrol—see Stilbestrol

Thyropoietin, not for average herd. Nov. 1956, p. 14

Weigh-a-day-a-month plan. Sept. 1956, p. 3

See also Dairy

Cecal worm and blackhead parasite. Aug. 1957, p. 12

Cheese, cheddar, new method. June 1956, p. 4; July 1956, p. 13

Chelates for fruit trees. June 1958, p. 16

Cherries, canned:

Cold storage. Dec. 1956, p. 16

Improved processing. Feb. 1958, p. 15

Chestnuts. Apr. 1957, p. 16; Jan. 1958, p. 15

## Chicken:

Breeding studies with fruit flies. July 1955, p. 11

Broiler production. Aug. 1955, p. 14; Aug. 1956, p. 5; Mar. 1958, p. 7

Brooder with curtains. Dec. 1956, p. 11

Cottonseed meal in feed. Nov. 1957, p. 15

Diseases—see Cecal worm, Chronic respiratory disease, Coccidiosis, Emergency measures

Feather meal. May 1956, p. 13

Frozen, fried, improving quality. Sept. 1955, p. 12

Management methods. July 1955, p. 4; May 1957, p. 4

More pounds on less feed. Sept. 1955, p. 10

## Chicken—Continued

National Poultry Improvement Plan. Dec. 1956, p. 15; Jan. 1958, p. 8

Nicarbazin, effect on egg production. June 1957, p. 15

Radioactive tracers. Oct. 1955, p. 13

Roasted, pink color in. Mar. 1956, p. 15

Selecting layers. July 1957, p. 4

Thyroid glands and production. Mar. 1956, p. 13

Waste disposal. Mar. 1957, p. 14

Children's height-weight handbook. Oct. 1957, p. 13

Chilling injury in stored fruits and vegetables. May 1957, p. 14

Chlorosis in soybeans. Feb. 1957, p. 11

Chromatography technique. Apr. 1956, p. 8

Chronic respiratory disease. Oct. 1956, p. 6; Sept. 1957, p. 15

## Citrus:

Nutrition. Aug. 1955, p. 5; May 1956, p. 10

Tristeza disease. Oct. 1955, p. 8

Cleaning cotton carpets—see Cotton

Cloth—see Fabric

## Clothing:

Choosing men's suits. Jan. 1958, p. 16

Will a T-shirt keep its shape? June 1958, p. 5

Cloud seeding to prevent forest fires. June 1957, p. 8

Clover—see Ladino clover, sweetclover

Coastal Bermuda grass. Dec. 1955, p. 16; Nov. 1956, p. 15

See also Grass

Cobalt 60—see Fruit flies, Screwworm

## Coccidiosis control:

Nicarbazin for chickens. June 1957, p. 15

Treatment for chicks. Nov. 1956, p. 15

Cockroaches for laboratory studies. July 1957, p. 8

Cocoa-butter substitute from fats. Dec. 1957, p. 14

## Color:

Comparator for judging wool value. Apr. 1958, p. 10

How sun makes apples red. Apr. 1957, p. 12

Keep canned cherries cool. Dec. 1956, p. 16

Conservation—see Soil, Water Management

## Consumers:

Benefit from agricultural research. Nov. 1957, p. 2

Facts for consumer education. Sept. 1957, p. 11

Contour farming. Oct. 1955, p. 7; June 1956, p. 16

Contract production of broils. Mar. 1958, p. 7

## Convenience foods:

City family survey. Mar. 1957, p. 8

Increasing demand for. Sept. 1956, p. 14

Cooking losses, beef. Mar. 1956, p. 15

Cooling produce. May 1956, p. 11; Aug. 1956, p. 12

Cooperatives. Sept. 1957, p. 16; Mar. 1958, p. 15; Apr. 1958, p. 15

Copper may harm citrus trees. May 1956, p. 10

Cork virus of sweetpotatoes, test for. June 1956, p. 11

## Corn:

Breeding earworm resistance. Nov. 1956, p. 3

Corn cob fractions make good roughage. Apr. 1958, p. 16

Dialdehyde starch for tanning. Dec. 1957, p. 5

High-amylose hybrids for industrial uses. Nov. 1957, p. 16

Irrigation. June 1956, p. 5; Dec. 1956, p. 7

New cash crop in Southeast spurs research. Oct. 1957, p. 5

Nitrogen for poor-land corn. Apr. 1956, p. 5



## Corn—Continued

- Predicting maturity for artificial drying. Sept. 1955, p. 7
- Ridge farming. Oct. 1955, p. 7
- Shelling saves space, money. Dec. 1956, p. 15
- Warm soil speeds growth. June 1957, p. 15

## Corn Borer:

- Granular insecticides show promise. Oct. 1955, p. 16
- 23-State survey. Mar. 1956, p. 7

Cortisone—*see Agave lecheguilla*; Yams, wild

Cost-price relationship in 1957. Dec. 1957, p. 3

Costs of production—*see Individual commodity*; Labor

## Cotton:

- Boll drop. May 1956, p. 6
  - Boll weevils resist insecticides. May 1957, p. 7
  - Breeding. Dec. 1956, p. 8; Apr. 1958, p. 6
  - Breeding for three-way disease resistance. Nov. 1955, p. 5
  - Carpets, cleaning. Feb. 1956, p. 15; Feb. 1957, p. 14
  - Coatings protect awnings. May 1957, p. 15; Mar. 1958, p. 16
  - Does defoliation pay? Aug. 1957, p. 15
  - Dye test shows fiber quality. Jan. 1956, p. 14
  - Fabric that sheds oil and water. June 1957, p. 15
  - Flame-resistance treatment for fabric. Oct. 1955, p. 16
  - Fully acetylated fabric. Oct. 1957, p. 15
  - Ginning. Sept. 1955, p. 8; Oct. 1956, p. 3; May 1957, p. 15; Jan. 1958, p. 14
  - How cotton soils. June 1958, p. 8
  - Insecticides, systemic. Aug. 1955, p. 16; Feb. 1957, p. 10
  - Pima S-1 variety. Mar. 1956, p. 3; May 1956, p. 16
  - Pink bollworm. July 1955, p. 6; Dec. 1955, p. 8; Feb. 1956, p. 16; Jan. 1958, p. 6
  - Radiation uses tested. Oct. 1956, p. 16; Aug. 1957, p. 16
  - Weed control. Apr. 1956, p. 5
- Cottonseed meal. Feb. 1957, p. 3; Nov. 1957, p. 15
- Cottonseed sorter, experimental. Mar. 1957, p. 16
- Coumestrol, new hormone from Ladino clover. Nov. 1957, p. 11
- CRD—*see Chronic respiratory disease*
- Cream, sweetened. Nov. 1957, p. 4
- Credit policies of cooperatives. Apr. 1958, p. 15
- Crooked snout—*see Atrophic rhinitis*

## Crops:

- Fertilize crop residues? Mar. 1956, p. 5
- Improvement through genetics. Mar. 1957, p. 2. *See also Individual crop*
- Regulatory organization, best defense against biological warfare. Oct. 1957, p. 6
- See also Plant introduction*

Crown rust—*see Rust*

Cucumbers, controlling pickle softening. Mar. 1957, p. 9

Cultural practices—*see Pink bollworm, Individual plant*

Culture collection. Apr. 1958, p. 8

Curing ham, home methods. Oct. 1955, p. 13

Curtains on chick brooders. Dec. 1956, p. 11

## D

## Dairy:

- Consumption of dairy products. June 1957, p. 10
- Dairy-hog farming, how to meet price drop. Dec. 1956, p. 6

## Dairy—Continued

Herd Improvement Association. Oct. 1955, p. 14; June 1956, p. 4; Sept. 1956, p. 3; Apr. 1958, p. 15

Herd Replacements. Feb. 1957, p. 5

Income increases on Northeast family farms. June 1956, p. 4

*See also Cattle, dairy*

Dalapon, how it kills weeds. Apr. 1958, p. 7

Defoliation, chemical, on cotton. Aug. 1957, p. 15

Dehydrated potato flakes. Nov. 1955, p. 12; Jan. 1957, p. 7

Dehydrofrozen peas. Dec. 1957, p. 15

## Detergents:

Animal fats in. Nov. 1956, p. 10

Choosing the right detergent. Dec. 1956, p. 13

Dialdehyde starch for tanning leather. Dec. 1957, p. 4

## Diet:

Daily food plan. Aug. 1956, p. 16; May 1958, p. 6

Fat—amount and source. Jan. 1958, p. 11

Older people need diet help. Feb. 1957, p. 13

*See also Food, Nutrition*

Diethyl toluamide repellent for mosquitoes. Mar. 1957, p. 6

Differentiator for sorting cottonseed. Mar. 1957, p. 16

Dioscorea—*see Yams, wild.*

## Disease:

In case of emergency. Aug. 1957, p. 3; Sept. 1957, p. 13; Oct. 1957, p. 2, 6

Need for research. Dec. 1955, p. 2

*See also Antibiotics, Virus, individual animal, plant, or disease.*

Distributor for air-spreading granular insecticides. Oct. 1957, p. 14

Diversification in Mississippi Delta. Sept. 1955, p. 6

Double cropping in South. Apr. 1956, p. 6

Dow ET-57, systemic for cattle grubs. July 1956, p. 15; Dec. 1956, p. 3; Jan. 1958, p. 15

DPA, herbicide helps rebuild pastures. Nov. 1955, p. 11

Drainage—*see Water management*

Droughts—*see Water management*

Dusting by plane. Sept. 1955, p. 14; Oct. 1957, p. 14

Dwarfs in cattle. Jan. 1958, p. 4

## Dye:

Making wool dye resistant. May 1957, p. 16

Test for cotton fiber quality. Jan. 1956, p. 14

## E

Earworm, breeding resistant corn. Nov. 1956, p. 3

Easter bulb storage. Nov. 1957, p. 14

## Economics:

Financing farm adjustments. Feb. 1957, p. 12

Future. June 1958, p. 2

How much to pay for land. Mar. 1958, p. 6

Living standards have improved greatly. Mar. 1958, p. 3

Looking back at 1957. Dec. 1957, p. 3

Our needs in 1975. Apr. 1957, p. 4

Price spread is increasing. Feb. 1958, p. 14

Puerto Rican development. Mar. 1958, p. 10

Ruralities in our expanding economy. Mar. 1957, p. 3

Tax appraisals, a way to improve. Jan. 1957, p. 5

Trends for 1958. Jan. 1958, p. 3



## Economics—Continued

What's keeping land prices up? Feb. 1956, p. 5

See also Chicken, Dairy, Fertilizer, Prices, Swine, individual commodity.

## Eggs:

Candling and egg quality. Jan. 1957, p. 11

Green-rot detector. Nov. 1956, p. 13

Membranes, whites hold back bacteria. Sept. 1957, p. 16

Elastin membrane and better leather. June 1957, p. 13

Electrical treatment for seeds. Jan. 1957, p. 3

Elevator for potatoes. Feb. 1957, p. 6

Emergency measures. Aug. 1957, p. 3; Sept. 1957, p. 13; Oct. 1957, p. 2, 6

Employment for ruralities. Mar. 1957, p. 3

Energy-saving kitchen—see House

Energy studies—see Metabolism

Equine infectious anemia—still unconquered. Oct. 1955, p. 11

## Equipment:

Home, service life. Feb. 1958, p. 10

Solar-energy collector for heating. Jan. 1956, p. 11

See also Housing, individual crop

Erosion—see Soil

## Estrogen:

Contaminated test feeds affect fertility. Apr. 1958, p. 11

Coumestrol from Ladino clover. Nov. 1957, p. 11

ET-57—see Dow ET-57

European chafer, new pesticide sought. Aug. 1957, p. 11

European corn borer—see Corn borer

Evapo-transpirometer. Dec. 1955, p. 6

Expenditures for family living. Mar. 1958, p. 3

Export certification for plants. Mar. 1958, p. 16

# F

## Fabric:

Facts for consumers. Sept. 1957, p. 11

Knit. May 1957, p. 11; June 1958, p. 5

See also Cotton, Wool

Facts for consumer education. Sept. 1957, p. 11

Fallout—see Radioactive

Family—see Economics, Food, Spending

## Farm:

Adjustments. Feb. 1957, p. 12

Fatal accidents. July 1957, p. 7

Planning can pay. Aug. 1955, p. 8

Trends in expanding economy. Mar. 1957, p. 3

See also Economics, Management

Farmers take vacations. Feb. 1957, p. 16

## Fat:

Carbohydrates, amino acids, and liver fat. Aug. 1956, p. 13

In human nutrition. May 1957, p. 12; Jan. 1958, p. 11; June 1958, p. 7

Measuring an animal's fat. Oct. 1956, p. 11

New uses for fats and oils. July 1955, p. 10

Substitute for imported cocoa butter. Dec. 1957, p. 14

Supplement in turkey diet. Sept. 1956, p. 13

Synthetic detergents with tallow. Nov. 1956, p. 10

Trimming from beef. Mar. 1956, p. 15

Use in candy. Nov. 1956, p. 16

Fatalities from farm accidents. July 1957, p. 7

Feather meal, growth factor in. May 1956, p. 13

## Feed:

Additives are widely used. Oct. 1956, p. 10

Efficiency. Jan. 1957, p. 2; Mar. 1957, p. 7. See also Individual animal

Estrogen contamination. Apr. 1958, p. 11

Fertility in turkeys—see Turkey, parthenogenesis

## Fertilizer:

Kit for fertilizer-use studies. May 1958, p. 15

Leaf analysis aids plant nutrition. Aug. 1955, p. 5

Pasture, use on. July 1955, p. 16; Dec. 1955, p. 16; Mar. 1956, p. 14; Nov. 1956, p. 15

Plant nutrition and insect control. June 1958, p. 4

Radioactive, for crop plant study. Feb. 1956, p. 8

Soil, moisture, nutrients, crops. May 1957, p. 6

See also Nitrogen, Phosphorus

## Fescue:

Making new grasses. Jan. 1956, p. 12

Winter pasturing and parasites in calves. June 1957, p. 6

Fever tick, cattle. Dec. 1957, p. 8

Filarial dermatosis in sheep. Mar. 1958, p. 13

Film, plastic, as reservoir liner. Dec. 1956, p. 10

Filtration-extraction process for vegetable oils. May 1956, p. 5

Financing farm adjustments. Feb. 1957, p. 12

Fire ant eradication effort. Sept. 1957, p. 15; Dec. 1957, p. 15

## Firefighting, forest:

Fighting by planes. Nov. 1957, p. 7

Skyfire project to combat fire-setting storms. June 1957, p. 8

Strip pastures as firebreaks. Oct. 1956, p. 6

Flakes, potato—see Potatoes

Flame-resistant cotton—see Cotton

Flaming reduces weed-control labor. Apr. 1956, p. 5

Flavor, single-sample taste test. June 1956, p. 15

Flax and disease. Dec. 1955, p. 3; May 1958, p. 16

Fleece—see Wool

Floods—see Water management

## Florida:

Range and cattle management. July 1957, p. 5

Water management. Nov. 1957, p. 12; Jan. 1958, p. 12

Flour beetle helps breeding studies. July 1955, p. 11

Flour quality. Mar. 1958, p. 4

Fluidizer may improve aerial dusting. Sept. 1955, p. 14

## Fly:

Attractant. Aug. 1955, p. 15

Insecticidal oil from heliopsis. Apr. 1957, p. 15

Shows promise for testing anticancer drugs. Apr. 1958, p. 14

## Food:

City families, food spending. Jan. 1957, p. 8

Convenience foods, increasing demand for. Sept. 1956, p. 14

Convenience foods, new versus old. Mar. 1957, p. 8

Daily Food Guide. Aug. 1956, p. 16; May 1958, p. 6

Dairy-products consumption. June 1957, p. 10

Data for nutritionists. Apr. 1956, p. 12

Losses in institutions. Sept. 1956, p. 14

Preservation in farm households. June 1956, p. 14

Tenderness in frozen poultry. Sept. 1956, p. 14

Use in households. May 1956, p. 15



## Food—Continued

*See also* Diet, Nutrition, individual food

Foot-and-mouth disease. Dec. 1955, p. 10; Sept. 1956, p. 7; Nov. 1957, p. 15

Forage—*see* Alfalfa, Grass, Pasture

## Forest:

Protecting seedlings from pests. Feb. 1957, p. 7

Timber resource review. Jan. 1956, p. 7

*See also* Fire, Gypsy moth, Trees

Fowl pox vaccination and parthenogenesis. Nov. 1957, p. 3

Freeze-dried potatoes for stock feed. Oct. 1955, p. 10

## Frozen food:

Home freezing on the farm. June 1956, p. 14

Lockers can expand services. Apr. 1957, p. 14

Testing, errors in. May 1958, p. 7

Time-temperature studies improve quality. Oct. 1957, p. 12

*See also* Individual food

## Fruit:

New varieties for northern Great Plains. Oct. 1957, p. 16

Precooling before packing. Aug. 1956, p. 12

*See also* Individual fruit

## Fruit flies:

In poultry breeding studies. July 1955, p. 11

Radiation, potential control weapon. Nov. 1957, p. 16;

Dec. 1957, p. 11

*See also* Mediterranean fruit fly

## Fruit trees:

Bark feeding. Sept. 1956, p. 11

Chelates for metal deficiencies. June 1958, p. 16

Fruit set, thinning sprays. Mar. 1956, p. 10; June 1957, p. 16

Fumigation—*see* Hall scale, Khapra beetle

## Fungi:

Aid trees. Mar. 1956, p. 6; June 1958, p. 3

Antibiotics for vegetables. Mar. 1956, p. 10; June 1956, p. 16

Camellia blight. Sept. 1957, p. 16

Carotene for vitamin A. Nov. 1957, p. 16

Culture collection. Apr. 1958, p. 8

Light's response to. July 1955, p. 12

Predicting late blight saves potatoes. Jan. 1956, p. 6

Ryegrass blind seed. May 1956, p. 4

*See also* Rust

## Fungicide:

Controls blackspot on roses. Mar. 1956, p. 16; Mar. 1957, p. 16

In tung oil varnish. June 1958, p. 6

Protects berry boxes. Oct. 1956, p. 15

Saving potatoes. Jan. 1956, p. 6

Fusarium wilt resistance in cotton. Nov. 1955, p. 5

# G

Garbage—*see* Vesicular exanthema

Genetics—*see* Breeding

Germination of hard seed controlled by temperature. Sept. 1955, p. 7

Gibberellic acid accelerates growth. Sept. 1956, p. 12

Ginning—*see* Cotton

Gossypol tolerance in swine related to protein. Feb. 1957, p. 3

Grapes, new varieties. Dec. 1955, p. 12; Oct. 1957, p. 15; May 1958, p. 11

## Grass:

African introductions. Feb. 1958, p. 6

Evaluating new forages. Sept. 1955, p. 2

For Mississippi Delta. Sept. 1955, p. 6

Making orchardgrass companionable. Oct. 1955, p. 2, 6

New varieties. Aug. 1955, p. 10; Jan. 1956, p. 12

Sericea lespedeza. Jan. 1957, p. 6

*See also* Fertilizer

## Grasshoppers:

Nutritional needs. Jan. 1958, p. 6

Threat in 1956. Feb. 1956, p. 6

Grazing—*see* Pasture, Range

Green-chop feed for dairy cattle. Oct. 1957, p. 15

Greenhouse, moisture control. Aug. 1956, p. 4

Green-rot detector for eggs. Nov. 1956, p. 13

Growth regulators—*see* Plant growth regulators

Growth standards for dairy cattle. Aug. 1955, p. 13

Grub, cattle. July 1956, p. 15; Dec. 1956, p. 3; Jan. 1958, p. 15; Mar. 1958, p. 12

Gypsy moth. May 1957, p. 8; June 1957, p. 7

# H

Hall scale. Jan. 1956, p. 5

Hallocks, fungicide prolongs life. Oct. 1956, p. 15

Halogeton. Apr. 1957, p. 11; May 1958, p. 15

Ham, home cured, keeping quality. Oct. 1955, p. 13

## Handling:

Apples. Sept. 1955, p. 13; Apr. 1957, p. 15

Potatoes. Feb. 1957, p. 6

Haploid cotton plants for breeding. Dec. 1956, p. 8

## Harvesting:

Apples. Sept. 1955, p. 13; Apr. 1957, p. 15

Beets. June 1956, p. 6

Castorbeans. Mar. 1957, p. 12

Potatoes. July 1956, p. 8

Prunes. June 1958, p. 14

Sugarcane. Sept. 1956, p. 5; Aug. 1957, p. 6

*See also* Seed, individual crop

Hatchability of turkey eggs. June 1956, p. 3

Hay mulch for weak apple trees. Jan. 1957, p. 14

Heat pump, help from sun. Jan. 1956, p. 11

Heat-tolerant cattle—*see* Cattle

Heat treatment for ratoon stunting in sugarcane. July 1955, p. 7

Heating stock water. Feb. 1957, p. 15

Heifer—*see* Cattle

Height-weight of children. Oct. 1957, p. 13

Heliopsis yields insecticidal oil. Apr. 1957, p. 15

Hendricks, Sterling. Apr. 1958, p. 3

Herbicides—*see* Pesticides, Weeds

Herd behavior and milk production. July 1955, p. 15

Hessian fly. Dec. 1955, p. 5; Feb. 1956, p. 7

Highway planting. Apr. 1957, p. 8

## Hog:

Aircraft noise. Jan. 1957, p. 10

Crossbreeding. Mar. 1957, p. 13



## Hog—Continued

- Diseases—see Atrophic rhinitis; Brucellosis, Parakeratosis; Vesicular exanthema
- Feed. Sept. 1955, p. 16; Oct. 1956, p. 10; Feb. 1957, p. 3
- Feeder-pig cost guide. Oct. 1957, p. 11
- Management in price drop. Dec. 1956, p. 6
- Semen ships well. Sept. 1957, p. 15
- Worms. Nov. 1956, p. 15; Feb. 1958, p. 13
- See also Livestock
- Hoja blanca rice disease. Dec. 1957, p. 15
- Holly. May 1957, p. 13
- Home economics research. Nov. 1955, p. 16; Sept. 1956, p. 16; Nov. 1956, p. 2
- See also Fabric, House, Nutrition
- Honey in baked goods. Mar. 1957, p. 15; see also Bees
- Horizontal silos with plastic. Feb. 1958, p. 12
- Hormone, coumestrol, from clover. Nov. 1957, p. 11
- Hormones in feed. Oct. 1956, p. 10; Apr. 1958, p. 11
- See also Stilbestrol
- Hornfly on dairy cattle, control. Apr. 1958, p. 15
- Horse—see Equine infectious anemia
- House:
  - Energy-saving kitchen. Oct. 1956, p. 8
  - Solar-energy for heating. Jan. 1956, p. 11
- Houseflies—see Fly
- Housing:
  - Cattle. Aug. 1957, p. 7
  - Chicken laying houses. May 1957, p. 4
  - Solar-energy collector. Jan. 1956, p. 11
- Huntley, Mont., Holsteins. Feb. 1956, p. 13
- Hydrangea color and aluminum. July 1957, p. 15
- Hydrocooling—see Carrots

## I

### Identification:

- Of insects. May 1957, p. 15
- Tags for cattle. Dec. 1955, p. 15
- Imported fire ant—see Fire ant
- Income—see Economics
- Increased use of agricultural products, commission on. Nov. 1956, p. 16
- Industrial products compete with agricultural. Oct. 1956, p. 2
- Insect:

- Beneficial lady beetle. Jan. 1956, p. 16
- Can plant nutrition affect insect control? June 1958, p. 4
- Identification. May 1957, p. 15
- Livestock pests. July 1956, p. 14
- Nematode against insects. Apr. 1956, p. 3
- Nutritional needs. Jan. 1958, p. 6
- Raising friendly insects. Aug. 1957, p. 9
- Repellent, all-purpose. Mar. 1957, p. 6
- Search for ally against halogeton. Apr. 1957, p. 11
- Surveys. Apr. 1958, p. 13
- Tree seedlings. Feb. 1957, p. 7
- Used for laboratory studies. July 1955, p. 11; July 1957, p. 8; Apr. 1958, p. 14
- See also Attractants, Insecticides, Nonchemical control, Pesticides, Plant Quarantine, individual insect

### Insecticides:

- Aerial spraying. Sept. 1955, p. 14; Feb. 1957, p. 16; Oct. 1957, p. 14
- Aerosol can, improved. Nov. 1957, p. 5
- Boosted by sesamol. Jan. 1956, p. 16
- Brine shrimp for residue tests. Oct. 1955, p. 16
- Cockroach for testing. July 1957, p. 8
- Controlling livestock pests. July 1956, p. 14
- For mosquito larvae. Sept. 1955, p. 16
- Granulars for corn borers. Oct. 1955, p. 16
- Lindane flavor in potatoes. July 1956, p. 10
- Needed: fruit moth spray that spares parasites. Aug. 1957, p. 11
- New safe, potent insecticides. Mar. 1958, p. 15
- Possible booster from sesame oil. Jan. 1956, p. 16
- Protecting stored peanuts. Sept. 1957, p. 6
- Residues in soil. Feb. 1958, p. 7
- Resistance to. July 1956, p. 16; Apr. 1957, p. 2; May 1957, p. 7, 16
- Respirator for all pesticides. Apr. 1957, p. 10
- Ryania as selective apple insecticide. Dec. 1955, p. 16
- Search for better insecticides. Jan. 1957, p. 13
- Sought: safe pesticide for nursery stock. Aug. 1957, p. 11
- Systemics for livestock. July 1956, p. 15; Dec. 1956, p. 3; Jan. 1958, p. 15; Mar. 1958, p. 12
- Systemics for plants. Aug. 1955, p. 16; Feb. 1957, p. 10; Jan. 1958, p. 16

### See also Nonchemical control, Pesticides

- Institution food losses and nutrition. Sept. 1956, p. 14
- Integration in poultry industry—see Chicken
- Interagency committee on nutrition—see Nutrition education
- Interdependence among scientific disciplines. May 1958, p. 2
- Introduction—see Nonchemical control, Plant introduction
- Investment:

- Cattle housing. Aug. 1957, p. 7
- Feeder pigs. Oct. 1957, p. 11
- Land. Mar. 1958, p. 6

- Iris blooms depend on storage temperatures. Nov. 1957, p. 14

### Irrigation:

- In pasture management. Sept. 1955, p. 6
- Of dry beans. Dec. 1956, p. 16
- Proper methods minimize mosquito breeding. Feb. 1957, p. 11
- Seeking irrigation answers. Dec. 1955, p. 6
- Timing for tobacco, corn. June 1956, p. 5
- Why irrigated corn lodges. Dec. 1956, p. 7
- See also Soil, Water management

- Isotopes—see Atomic energy, Radioactive

## J

- Japanese beetle, new pesticide sought. Aug. 1957, p. 11
- Jelly. Dec. 1957, p. 14; Apr. 1958, p. 16
- Johne's disease. Apr. 1958, p. 11
- Johnsongrass controlled by dalapon. Apr. 1958, p. 7

## K

- Khapra beetle fumigation. July 1955, p. 16; Mar. 1956, p. 8
- Kitchen—see Housing



Klamath weed destroyed by beetle. July 1955, p. 5  
Knit fabric—see Fabric  
Knot-free wood from pruned pine. Oct. 1956, p. 16

## L

Labor. July 1955, p. 4; May 1957, p. 4; Nov. 1957, p. 8  
Labor saving—see Harvesting. Labor. Particular product  
Laboratories, USDA:  
    Animal. Aug. 1955, p. 14; Sept. 1956, p. 7, 16; Oct. 1956, p. 16; Mar. 1957, p. 15; July 1957, p. 2  
    Crops, seeds, soils. Sept. 1957, p. 3; Nov. 1957, p. 15  
    Pioneering. Aug. 1957, p. 2; Jan. 1958, p. 2; June 1958, p. 15  
    Utilization. Apr. 1956, p. 2; Apr. 1957, p. 3  
Lactometer to measure milk solids. Nov. 1957, p. 6  
Ladino clover:  
    Source of estrogen. Nov. 1957, p. 11  
    Virus infection. Apr. 1956, p. 7; Feb. 1958, p. 16  
Lambs—see Sheep  
Land:  
    Management. July 1955, p. 8; Aug. 1955, p. 8; Mar. 1958, p. 10  
    Values. Feb. 1956, p. 5; Jan. 1957, p. 5; Dec. 1957, p. 3; Mar. 1958, p. 6  
    See also Management  
Laundering detergents. Dec. 1956, p. 13  
Leaf:  
    Analysis helps solve nutritional problems. Aug. 1955, p. 5  
    Leaf-rust resistance in wheat. Aug. 1956, p. 8  
Leather:  
    Dialdehyde starch for tanning. Dec. 1957, p. 4  
    Seeking ways to modify elastin membrane. June 1957, p. 13  
Legumes, weedkillers for. Mar. 1956, p. 5; June 1957, p. 15  
Light:  
    Effect on turkey fertility. Oct. 1957, p. 10  
    How sun makes apples red. Apr. 1957, p. 12  
    Regulating *Weigela* plant growth. May 1956, p. 16  
    Response of fungi. July 1955, p. 12  
    What day length means to trees. Dec. 1956, p. 10  
Lightning—see Skyfire  
Lilies and storage temperatures. Nov. 1957, p. 14  
Lima beans:  
    New nematode-resistant variety. Apr. 1958, p. 5  
    Streptomycin controls downy mildew. Mar. 1956, p. 10  
Lime—see Tristeza  
Lindane—see Insecticides  
Linoleic acid—see Fat  
Lipids in flour and baking quality. Mar. 1958, p. 4  
Livestock:  
    Aircraft noise. Jan. 1957, p. 10  
    Auctions, improving. Nov. 1956, p. 11  
    Costs of producing. July 1955, p. 2, 4  
    Diseases. Aug. 1955, p. 12; Dec. 1955, p. 2; Sept. 1957, p. 13  
    Feed additives. Oct. 1956, p. 10  
    Insect control. July 1956, p. 14; see also Nonchemical control, Pesticides, Systemics  
    Packing operation may pay. Oct. 1957, p. 11  
    Poisoning from pesticides. May 1958, p. 4

## Livestock—Continued

    Warming stock water. Feb. 1957, p. 15  
    See also Cattle, Chickens, Hogs, Sheep  
Living standards of farm families. Mar. 1958, p. 3  
Locker plants can expand. Apr. 1957, p. 14  
Lodging—see Corn  
Logs—see Timber  
Lungworm. Dec. 1955, p. 11  
Lupine, Borre sweet blue. Dec. 1957, p. 4  
Lure—see Attractant  
Lysimeter in water studies. Apr. 1957, p. 6

## M

Magnesium for apples. Jan. 1957, p. 14  
Management, farm. Aug. 1955, p. 8; Dec. 1956, p. 6; Mar. 1957, p. 10; Mar. 1958, p. 10  
Manganese may harm citrus trees. May 1956, p. 10  
Maple flavor boosted by bacteria. Feb. 1957, p. 15  
Marketing. Sept. 1956, p. 4; Sept. 1957, p. 16; Feb. 1958, p. 14  
    See also Auctions  
Maturity in corn, predicting. Sept. 1955, p. 7  
Meat:  
    Breeding for tenderness. Feb. 1957, p. 15  
    Inspection. June 1956, p. 7; July 1957, p. 10; Mar. 1958, p. 15  
    Packing operation may pay. Oct. 1957, p. 11  
    See also Beef, Ham, Pork, Poultry  
Mechanization—see Double-cropping, Harvesting, particular crop  
Mediterranean fruit fly. July 1956, p. 10; Nov. 1956, p. 5; Apr. 1957, p. 13; July 1957, p. 16; Sept. 1957, p. 7  
Men's suits. Jan. 1958, p. 16  
Metabolism:  
    Dairy cow. May 1958, p. 8; June 1958, p. 13  
    Pioneering laboratory in cellular metabolism. June 1958, p. 15  
Methoxychlor dust for hornfly. Apr. 1958, p. 15  
Micronutrients:  
    Abnormal soils. Nov. 1955, p. 6  
    Bark feeding fruit trees. Sept. 1956, p. 11  
    Food-quality frontier. Oct. 1955, p. 3  
    Overfeeding citrus trees. May 1956, p. 10  
Micro-organism collection. Apr. 1958, p. 8  
Mildew, powdery, in apples. Mar. 1956, p. 11  
Military preparedness and research. June 1956, p. 2  
Milk:  
    Cost of home produced. June 1957, p. 10  
    Drying processes for nonfat solids. Apr. 1956, p. 10  
    Marketing margins. Sept. 1956, p. 4  
    Nonfat solids test kit. Nov. 1957, p. 6  
    Production records—see Dairy Herd Improvement Assn.  
    Vending machines. Jan. 1957, p. 16  
    Vitamin B<sub>12</sub>. Oct. 1956, p. 12  
    See also Cattle, dairy; Dairy  
Minerals, Trace—see Micronutrients  
Minks, vaccine for virus enteritis. June 1958, p. 12  
Mitochondria in sweetpotatoes isolated. May 1957, p. 14  
Moisture control in greenhouse. Aug. 1956, p. 4  
Molasses for livestock. Dec. 1956, p. 5



Mold causes pickle softening. Mar. 1957, p. 9

MOPA:

- Growth regulator. Feb. 1958, p. 12
- Weapon for nematodes. Feb. 1956, p. 12; June 1956, p. 12

Mosquitoes:

- New repellent. Mar. 1957, p. 6
- Proper irrigation minimizes breeding. Feb. 1957, p. 11
- Resistance to insecticides. May 1957, p. 16

Muck soils, conserving. Oct. 1957, p. 3

Mulch:

- Enriched hay revives apple trees. Jan. 1957, p. 14
- In water conservation. Apr. 1957, p. 6
- Related to lower corn yields. June 1957, p. 15
- Winter wheat—stubble beats drought. Feb. 1957, p. 15

Mushrooms:

- Chlorinated water spray. Mar. 1956, p. 11
- Improvements through research. Mar. 1958, p. 8

Mycorhiza—see Fungi

## N

NAA fruit thinning spray improved. June 1957, p. 16

National Poultry and Turkey Improvement Plans—see Chicken

National Seed Storage Laboratory. Sept. 1956, p. 6

Naval stores—see Pine gum, Rosin

Nematodes:

- Against insects. Apr. 1956, p. 3
- Golden, from England. Sept. 1957, p. 15
- Pine seedlings. Feb. 1957, p. 7
- Root-knot. Nov. 1955, p. 5; Apr. 1956, p. 16; June 1956, p. 12; Apr. 1958, p. 5
- Search for nematocides. Feb. 1955, p. 6
- See also Soybeans

Neutron counter measures soil moisture. Dec. 1955, p. 6

Nicarbazin lowers egg production. June 1957, p. 15

Nitrogen:

- Can increase lodging in corn. Dec. 1956, p. 7
- Cattle like enriched grass. Nov. 1956, p. 15
- Enriched hay mulch revives apples. Jan. 1957, p. 14
- Fertilize crop residues? Mar. 1956, p. 5
- For poor-land corn. Apr. 1956, p. 5
- Low-cost applicator for liquid N. Nov. 1955, p. 10
- Needed for canned-peach quality. Apr. 1957, p. 12

Noise, aircraft, and livestock. Jan. 1957, p. 10

Nonchemical control:

- Attractants. Aug. 1955, p. 15; June 1956, p. 12; May 1957, p. 8; June 1957, p. 7; Sept. 1957, p. 7, 15; see also Mediterranean fruit fly
- Beetle controls Klamath weed. July 1955, p. 5
- Beneficial lady beetle. Jan. 1956, p. 16
- Blind-seed disease. May 1956, p. 4
- Breeding—see Breeding resistant varieties
- Editorial. Apr. 1957, p. 2
- Fescue pasture and parasites. June 1957, p. 6
- Flaming, spray for cotton weeds. Apr. 1956, p. 5
- Fungi aid trees. Mar. 1956, p. 6; June 1958, p. 3
- Halogeton. Apr. 1957, p. 11; May 1958, p. 15
- Hessian fly. Dec. 1955, p. 5; Feb. 1956, p. 7
- Hog pastures and whipworms. Nov. 1956, p. 15
- Nematode against insects. Apr. 1956, p. 3

## Nonchemical control—Continued

- Nematode repellent in tomatoes. June 1956, p. 12
- Nutritional needs of insects. Jan. 1958, p. 6
- Oriental fruit moth parasites. July 1957, p. 16; Aug. 1957, p. 11
- Pine seedlings. Feb. 1957, p. 7
- Pink bollworm. July 1955, p. 6; Dec. 1955, p. 8; Feb. 1956, p. 16; Jan. 1958, p. 6
- Plant nutrition and insect control. June 1958, p. 4
- Raising friendly insects. Aug. 1957, p. 8
- Shock treatment for weed seeds. Jan. 1957, p. 3
- Sterile male technique. (Fruit flies) Nov. 1957, p. 16; Dec. 1957, p. 11; (Screwworm) Nov. 1955, p. 3; July 1957, p. 15; Dec. 1957, p. 15
- Tapeworm and sanitation. July 1957, p. 10
- Virus against sawflies. Apr. 1956, p. 16
- Virus combats tent caterpillar. Dec. 1957, p. 7
- Wasp ally for tobacco. July 1956, p. 3
- See also Emergency measures, Plant quarantine, Regulatory Work

Nursery stock and quarantines. Aug. 1957, p. 11

Nutrition:

- Carbohydrates, amino acids, and liver fat. Aug. 1956, p. 13
- Effect of institution food loss. Sept. 1956, p. 14
- Frontier studies in plant, animal nutrition. Oct. 1955, p. 3
- Need for consumer education. Dec. 1956, p. 14
- Older people need diet help. Feb. 1957, p. 13
- Pioneering Laboratory in Cellular Metabolism. June 1958, p. 15
- Research needs. Jan. 1958, p. 16
- Survey of U.S. eating habits. May 1957, p. 12
- See also Fat, Food, Pantothenic acid, Vitamins, particular nutrient

Nutrition, plant—see Plant nutrition

## O

Oats:

- Crown rust. May 1956, p. 3
- New bronco variety. Feb. 1957, p. 16
- We're ahead of oat diseases. July 1956, p. 6

Off-flavor—see Flavor

Ohio station jubilee. Dec. 1957, p. 15

Oilseeds, new filtration-extraction process. May 1956, p. 5

Oil—see Fat

Orange—see Citrus

Orchardgrass, breeding. Oct. 1955, p. 6

Oriental fruit moth parasites. July 1957, p. 16; Aug. 1957, p. 11

Oriental lady beetle for pests. Jan. 1956, p. 16

Ornamentals. July 1956, p. 12; Dec. 1956, p. 15; May 1957, p. 16; July 1957, p. 14

## P

Palustric acid, new chemical from rosin. Dec. 1955, p. 14

Pantothenic acid assay method. Nov. 1956, p. 8

Parakeratosis cured by zinc. Apr. 1957, p. 5

Parasites:

- Anaplasmosis. May 1957, p. 3
- Beetle for Klamath Weed. July 1955, p. 5



## Parasites—Continued

- Blackhead and cecal worm. Aug. 1957, p. 12
- Cattle fever tick. Dec. 1957, p. 8
- Cattle grub. July 1956, p. 15; Dec. 1956, p. 3; Jan. 1958, p. 15; Mar. 1958, p. 12
- Filarial dermatosis. Mar. 1958, p. 13
- For insect control. July 1957, p. 16; Aug. 1957, p. 9, 11
- Grub systemic for roundworms. Jan. 1958, p. 15
- Lungworm immunity. Dec. 1955, p. 11
- Tapeworm. July 1957, p. 10
- Winter pasture and worms. June 1957, p. 6
- Witchweed. Jan. 1957, p. 15; Apr. 1958, p. 16
- Worms in hogs. Nov. 1956, p. 15; Feb. 1958, p. 13
- See also* Biological warfare defense, Fungi, Nematode, Screwworm

## Parthenogenesis—see Turkey

## Pasture:

- Alfalfa can get along with grass. Nov. 1955, p. 11
- All-season forage for Delta region. Sept. 1955, p. 6
- Enough fertilizer to pay. Mar. 1956, p. 14
- Herbicides can help rebuild pastures. Nov. 1955, p. 11
- Lupine may give South year-round grazing. Dec. 1957, p. 4
- Management. Mar. 1957, p. 10
- Parasites and winter pasture. June 1957, p. 6
- Pasture research, cooperation. Oct. 1955, p. 2
- Strip pastures fight fires. Oct. 1956, p. 6
- What's good for the Blacklands? July 1955, p. 8
- See also* Alfalfa, Grass

## Peaches:

- Mite carries mosaic. Sept. 1955, p. 15
- Needed: fruit moth spray that spares parasites. Aug. 1957, p. 11
- New concentrates. July 1957, p. 15
- Nitrogen for canned-peach quality. Apr. 1957, p. 12
- Training trees. May 1958, p. 12

## Peanuts, stored, protecting from insects. Sept. 1957, p. 6

## Pears:

- Canning waste for feed. Apr. 1957, p. 15
- Fruit-set spray. Mar. 1956, p. 10
- Ripening in transit. June 1956, p. 13
- Thinning. Dec. 1956, p. 15

## Peas, dehydrofrozen. Dec. 1957, p. 15

## Pepper, black, for Puerto Rico crop. July 1955, p. 14

## Personnel, USDA:

- Aug. 1955, p. 16; Nov. 1955, p. 16; Jan. 1956, p. 16; Mar. 1956, p. 16; Aug. 1956, p. 16; Sept. 1956, p. 16; May 1957, p. 2; May 1958, p. 16
- See also* Awards

## Pest control—see Nonchemical control

## Pesticides:

- Pests—see Biological warfare, Eradication, Insects, Nematodes, Nonchemical control, Regulatory work, Trees
- Livestock. July 1956, p. 14
- One respirator for all. Apr. 1957, p. 10
- Poisoning in livestock. May 1958, p. 4
- See also* Fungicide, Insecticides, Nematode, Systemics, Weeds

## Phosphorus nutrition of plants. Feb. 1956, p. 10

## Photosynthesis. Nov. 1955, p. 2

## Picking boxes for fruit—see Hallocks

## Pickle softening, control. Mar. 1957, p. 12

## Pigment:

- How sun reddens apples. Apr. 1957, p. 12
- Coating for awnings. Mar. 1958, p. 16

## Pima—see Cotton

## Pine:

- Gum derivatives for synthetic rubber. Mar. 1958, p. 15
- Loblolly, thinning. June 1956, p. 16
- Protecting seedlings from pests. Feb. 1957, p. 7
- Pruning prevents knots in wood. Oct. 1956, p. 16
- Sawfly control. Apr. 1956, p. 16

## Pink bollworm:

- Cultural control practices. July 1955, p. 6
- Nutritional needs. Jan. 1958, p. 6
- Wild-cotton hosts. Dec. 1955, p. 8; Feb. 1956; p. 16

## Pioneering laboratories—see Laboratories, USDA

## Piperazines for worms in hogs. Feb. 1958, p. 13

## Piroplasmosis. Dec. 1957, p. 8

## Pistaschio nuts, new crop. Oct. 1956, p. 13

## Plane—see Aircraft

## Plant cells, cracking. May 1957, p. 14

## Plant diseases:

- Antibiotics for vegetables. Mar. 1956, p. 10; June 1956, p. 16
- Tests for. June 1956, p. 11; Oct. 1956, p. 5; Feb. 1957, p. 8
- See also* Breeding resistant varieties, Fungi, Nonchemical control, Virus, individual plant or disease

## Plant export certification. Mar. 1958, p. 16

## Planting equipment. Apr. 1956, p. 6

## Planting for roadside cover. Apr. 1957, p. 8

## Plant growth regulators:

- Fruit set spray for pears. Mar. 1956, p. 10
- Gibberellic acid. Sept. 1956, p. 12
- Influence on boll drop. May 1956, p. 6
- Interplant transfer. Feb. 1958, p. 12
- Potato sprout inhibitor. Sept. 1955, p. 16

## Plant introduction:

- And seed bank. Sept. 1956, p. 6
- Blackberries. June 1957, p. 3
- Black pepper. July 1955, p. 14
- Developing new crops. Dec. 1957, p. 2
- Grasses from Africa. Feb. 1958, p. 6
- Ornamentals. July 1956, p. 12; Dec. 1956, p. 15; May 1957, p. 13, 16; July 1957, p. 14
- Pistaschio nuts. Oct. 1956, p. 13
- Strawberries. June 1957, p. 3
- Sugarcane. Nov. 1957, p. 10
- Yams for cortisone. Jan. 1956, p. 8; Feb. 1958, p. 15

## Plant nutrition:

- Effect of soil moisture. May 1957, p. 6
- Frontier studies. Oct. 1955, p. 3
- How plants get phosphorus. Feb. 1956, p. 10
- Leaf analysis. Aug. 1955, p. 5
- May affect insect control. June 1958, p. 4
- Micronutrient-abnormal soils. Nov. 1955, p. 6
- See also* Fertilizer, Fruit trees, Micronutrients



Plant pest control—see Fungicide, Insecticides, Insects, Nematode, Pesticides, Pests, Plant quarantine, Regulatory work, Weedkillers

Plant, photosynthesis. Nov. 1955, p. 2

Plant quarantine. Aug. 1956, p. 14; Aug. 1957, p. 11; Feb. 1958, p. 8; see also Soybean cyst nematode

Plastic film:

- As reservoir liner. Dec. 1956, p. 10
- For horizontal silos. Feb. 1958, p. 12
- Packaged carrots. May 1956, p. 11

Plowing, contour. June 1956, p. 16

Plum Island—see Foot-and-mouth disease; Laboratories, USDA

Poison—see Toxicity

*Polistes exclamans*. July 1956, p. 3

Polyploid—see Grapes, Grass

Pomace, apple, for cattle. May 1958, p. 3

Pork, rancidity in frozen. Jan. 1958, p. 10

Potatoes:

- Blight protection. Jan. 1956, p. 6
- Elevator for handling. Feb. 1957, p. 6
- Facts for consumers. Sept. 1957, p. 11
- Flakes. Nov. 1955, p. 12; Jan. 1957, p. 7
- Freeze-dried for cattle. Oct. 1955, p. 10
- French fries. Apr. 1956, p. 11
- Harvesting mechanically. July 1956, p. 8
- Improved varieties. Nov. 1955, p. 13
- Off-flavor from lindane. July 1956, p. 10
- Prepeeled. May 1957, p. 10
- Scab resistance. Aug. 1955, p. 16
- Sprout inhibitor. Sept. 1955, p. 16
- Thiodan controls green peach aphid. June 1958, p. 15
- Verticillium wilt resistance. July 1955, p. 13
- Wireworm now resists chlordane. Jan. 1957, p. 13

Poultry:

- Frozen, processing for tenderness. Sept. 1956, p. 14
- Waste disposal. March 1957, p. 14
- See also Chicken, Turkey

Precooling—see Cooling

Predators—see Nonchemical control

Prepeeled potatoes. May 1957, p. 10

Preservation, food—see Canning, Dehydrofrozen, Frozen, Jelly

Price:

- Drop, dairy-hog, how to meet. Dec. 1956, p. 6
- In 1957. Dec. 1957, p. 3
- Land. Feb. 1956, p. 5; Mar. 1958, p. 6
- Spread. Feb. 1958, p. 14
- Trends for 1958. Jan. 1958, p. 3
- See also Economics

Processed foods—see Food, Jelly, Potatoes

Production:

- Costs—see Labor, individual commodity
- Milk—see Dairy Herd Improvement Assn., Herd behavior
- Needs in 1975. April 1957, p. 4

Protein:

- Low-protein diet may hurt calf crop. Apr. 1956, p. 13
- Measuring, two easy ways. Dec. 1957, p. 6
- Needs of beef calves. July 1957, p. 12
- Versus gossypol in cottonseed meal. Feb. 1957, p. 3
- See also Amino acids

Prune harvester. June 1958, p. 14

Psorosis, citrus virus disease. Oct. 1955, p. 9

Puerto Rico:

- Growing black pepper. July 1955, p. 14
- Improving economy. Mar. 1958, p. 10

Pulpwood from sawmill wastes. May 1958, p. 14

## Q

Quarantine—see Brucellosis, Foot-and-mouth disease, Plant Quarantine, Soybean cyst nematode

## R

Rabbits produce antibodies for crop-disease work. Feb. 1956, p. 3

Radiation—see Cotton, Fruit flies, Screwworm

Radioactive fallout. Aug. 1957, p. 3, 16; Sept. 1957, p. 13: see also Atomic energy

Radiometer in irrigation. Dec. 1955, p. 6

Rancidity in frozen pork. Jan. 1958, p. 10

Range management:

- It pays to graze moderately. May 1956, p. 12
- More from Florida range. July 1957, p. 5
- See also Pasture

Ratoon stunting disease. July 1955, p. 7; Aug. 1956, p. 10

Rats may carry atrophic rhinitis. Sept. 1957, p. 12

Record of performance—see particular animal

Red color in apples. Apr. 1957, p. 12

Refrigeration of pears in transit. June 1956, p. 13

Regulatory work. Jan. 1956, p. 2. See also Emergency measures, Plant quarantine, Meat inspection

Repellant, insect, all-purpose. Mar. 1957, p. 6

Repellant in tomato roots. June 1956, p. 12

Reproduction in cattle and protein. Apr. 1956, p. 13

Research:

- Aids military preparedness. June 1956, p. 2
- Basic. Dec. 1956, p. 2; Feb. 1958, p. 3
- Confidence in results. Feb. 1958, p. 2
- Coordination, is more needed? Feb. 1956, p. 2
- Importance. May 1956, p. 2
- Many sciences. Aug. 1956, p. 2
- Need for. June 1957, p. 2
- On farm-product use. Apr. 1957, p. 3
- Value of. Sept. 1957, p. 2

Reservoir liner, plastic. Dec. 1956, p. 10

Residues:

- Fertilizer crop residues? Mar. 1956, p. 5
- Insecticide, in soil. Feb. 1958, p. 7

Respirator for all pesticides. Apr. 1957, p. 10

Rice:

- Drying. June 1958, p. 15
- Hoja blanca disease. Dec. 1957, p. 15
- Quality testing. Aug. 1957, p. 5

Ridge farming solves problems. Oct. 1955, p. 7

Roach—see Cockroach

Roadside cover for highways. Apr. 1957, p. 8

Root-knot nematode:

- Lima beans, resistant variety. Apr. 1958, p. 5
- On tomatoes. June 1956, p. 12
- Resistance in cotton. Nov. 1955, p. 5
- Soil texture. Apr. 1956, p. 16



ROP (record of performance)—*see* particular animal  
 Roses, blackspot control. Mar. 1956, p. 16; Mar. 1957, p. 16  
 Rosin, yields new acid. Dec. 1955, p. 14  
 Rotation grazing plan for delta region. Sept. 1955, p. 6  
 Rubber:  
   Natural rubber synthesized. Jan. 1957, p. 15  
   Pine-gum derivatives. Mar. 1958, p. 15  
 Rumen:  
   Bacteria in sheep. Aug. 1955, p. 11  
   Calves don't need inoculation. June 1957, p. 11  
 Ruralites in expanding economy. Mar. 1957, p. 3  
 Rust:  
   Flax. Dec. 1955, p. 3  
   Oat. May 1965, p. 3; July 1956, p. 6  
 Rustleometer for measuring handle in fabric. Aug. 1956, p. 6  
 Ryania insecticide for apples. Dec. 1955, p. 16  
 Ryegrass, blind seed disease curbed. May 1956, p. 4

## S

Sagebrush, potential gain from control. May 1958, p. 15  
 Sanitation necessary to stop tapeworm. July 1957, p. 10  
 Sawflies on pine killed by virus. Apr. 1956, p. 16  
 Sawmill wastes, savaging. May 1958, p. 14  
 School lunch based on research. Dec. 1957, p. 12  
 Screwworm:  
   Spray. Mar. 1958, p. 12  
   Sterile male technique. Nov. 1955, p. 3; July 1957, p. 15;  
   Dec. 1957, p. 15  
 Seed:  
   Big seeds for range. May 1957, p. 15  
   Blind-seed ryegrass disease curbed. May 1956, p. 4  
   Effect of temperature on sprouting. Sept. 1955, p. 7  
   Harvest device for shattered seed. Oct. 1956, p. 16  
   Seed storage laboratory. Sept. 1956, p. 6  
   Shock treatment. Jan. 1957, p. 3  
   Treated with insecticide. Feb. 1957, p. 10  
 Seedlings, foresters are saving. Feb. 1957, p. 6  
 Sericea lespedeza may be improved. Jan. 1957, p. 6  
 Sesame, improved varieties. June 1957, p. 4  
 Sesamolins may improve insecticides. Jan. 1956, p. 16  
 Sewing, pin or thread baste? July 1957, p. 16  
 Shade for livestock. Oct. 1957, p. 16  
 Sheep:  
   Breeding for wool. Sept. 1955, p. 3  
   Food from farm wastes. Aug. 1955, p. 11  
   Hybrid lambs outclass purebreds. Jan. 1956, p. 10  
   Open-faced, smooth sheep for range. Mar. 1956, p. 12  
   Rumen injections combat "sore head." Mar. 1958, p. 13  
   Selection for breeding. June 1957, p. 12  
   *See also* Livestock, Wool  
 Shelterbelts—*see* Windbreaks  
 Shirts, knit, and shape holding. June 1958, p. 5  
 Shock treatment for seeds. Jan. 1957, p. 3  
 Shrink-resistant wool. July 1957, p. 6  
 Sick time of farmers. Mar. 1957, p. 16  
 Silage:  
   Bacteria affect quality. Aug. 1957, p. 10  
   Poisonous gas in. Jan. 1957, p. 16

## Silage—Continued

  Production increasing. June 1958, p. 12  
   Trench silos. Dec. 1957, p. 16; Feb. 1958, p. 12  
 Size of children. Oct. 1957, p. 13  
 Skyfire studies to prevent forest fires. June 1957, p. 8  
 Slingshot sorter for cotton seed. Mar. 1957, p. 16  
 Sodium orthophenyl phenate protects berry boxes. Oct. 1956,  
   p. 15  
 Soil:  
   Conservation and management. July 1956, p. 2  
   Conserving muck soils. Oct. 1957, p. 3  
   Cover crops and soil crumb. Jan. 1958, p. 15  
   Deep tillage, when and how. Sept. 1957, p. 4  
   Frontier studies in nutrition. Oct. 1955, p. 3  
   How plants get phosphorus. Feb. 1956, p. 10  
   Micronutrient-abnormal soils, search for. Nov. 1955, p. 6  
   Moisture, nutrients, and crops. May 1957, p. 6  
   Mulch related to lower corn yields. June 1957, p. 15  
   Parallel terraces pay. June 1956, p. 5  
   Ridge farming solves problems. Oct. 1955, p. 7  
   Samples, foil containers for. Mar. 1957, p. 15  
   Seeking irrigation answers. Dec. 1955, p. 6  
   Tax appraisal on earning power. Jan. 1957, p. 5  
   What's good for the Blacklands? July 1955, p. 8  
   Yearbook on soil. Oct. 1957, p. 4  
   *See also* Fertilizer, Water management  
 Soil-resistant finish for cotton. June 1958, p. 8  
 Solar energy for heat pump. Jan. 1956, p. 11  
 "Sore head" in sheep—*see* Filarial dermatosis  
 Sorghum:  
   Breeding. May 1956, p. 16  
   New sirup sorgho. Aug. 1956, p. 11  
   Potential sugar crop. June 1958, p. 16  
   Promising sorghum-grass crop for South. Aug. 1955, p. 10  
   Stored moisture for grain sorghum. May 1957, p. 5  
 Soybeans:  
   Breeding for nematode resistance. Feb. 1958, p. 15  
   Double cropping in South. Apr. 1956, p. 6  
   Grant variety for North. Dec. 1955, p. 5  
   Is chlorosis bad? Feb. 1957, p. 11  
   Soybean cyst nematode. Feb. 1956, p. 16; June 1957, p. 16;  
   Aug. 1957, p. 16; Dec. 1957, p. 16  
 Spending:  
   Farm families. Mar. 1958, p. 3  
   For food by city families. Jan. 1957, p. 8; Mar. 1957, p. 8  
 Spider mites now resist insecticides. July 1956, p. 16  
 Spotted alfalfa aphid is spreading. July 1956, p. 16  
 Spraying by plane—*see* Aircraft  
 Starch, dialdehyde, for tanning. Dec. 1957, p. 4  
 State experiment station cooperation. Oct. 1955, p. 2  
 Statistics summer session. Apr. 1957, p. 15  
 Sterile male technique—*see* Fruit flies, screwworm  
 Sterility from contaminated feeds. Apr. 1958, p. 11  
 Stilbestrol. Nov. 1955, p. 16; Oct. 1956, p. 12; Mar. 1956, p.  
   16; June 1958, p. 15  
 Storage:  
   Effect on beans. Dec. 1956, p. 12  
   Food—*see* Individual foods  
   Peanuts. Sept. 1957, p. 6  
   Temperatures for bulbs. Nov. 1957, p. 14  
 Strawberries from South America. June 1957, p. 3

- Streptomycin—*see* Antibiotics
- Stripe-mosaic virus, new infection tests. Feb. 1957, p. 8
- Stubble-mulched wheat beats drought. Feb. 1957, p. 15
- Sugarbeets, mechanizing production. June 1956, p. 6
- Sugarcane:
- Bagasse, new uses. July 1956, p. 4; June 1958, p. 16
  - Fresh is worth more. Sept. 1956, p. 5
  - Introductions. Nov. 1957, p. 10
  - New harvester defoliates cane. Aug. 1957, p. 6
  - Ratoon stunting disease. July 1955, p. 7; Aug. 1956, p. 10
  - Wax obtained from cosmetic cane. May 1958, p. 10
- Suits, men's judging quality. Jan. 1958, p. 16
- Sun-energy collector for heating. Jan. 1956, p. 11
- Surveys:
- Accidents, fatal. July 1957, p. 7
  - Broiler production. Mar. 1958, p. 7
  - Children's height-weight data. Oct. 1957, p. 13
  - City families, food spending. Jan. 1957, p. 8
  - Convenience foods, use of. Sept. 1956, p. 14; Mar. 1957, p. 8
  - Corn borers. Mar. 1956, p. 7
  - Dairy product consumption. June 1957, p. 10
  - Family farm income. June 1956, p. 4
  - Farmer vacations. Feb. 1957, p. 16
  - Farm living standards. Mar. 1958, p. 3
  - Food use. May 1956, p. 15; May 1957, p. 12
  - Home canning. June 1956, p. 14
  - Labor in livestock production. July 1955, p. 4
  - Land prices. Feb. 1956, p. 5
  - Marketing. Sept. 1956, p. 4; Sept. 1957, p. 16
  - Price spread. Feb. 1958, p. 14
  - Ruralites in expanding economy. Mar. 1957, p. 3
  - Sick time. Mar. 1957, p. 16
  - Silage production. June 1958, p. 12
  - Timber Resource Review. Jan. 1956, p. 7
  - Trends in 1957. Dec. 1957, p. 3
- Swamp fever—*see* Equine infectious anemia
- Sweetclover on Texas Blacklands. July 1955, p. 8
- Sweetpotatoes:
- Cell studies for improving food quality. May 1957, p. 14
  - New products. Aug. 1957, p. 15
  - Test to detect cork virus. June 1956, p. 11
- Symbiosis—*see* Fungi
- Synthetic detergents—*see* Detergents
- Systemics:
- Livestock. July 1956, p. 15; Dec. 1956, p. 3; Jan. 1958, p. 15; Mar. 1958, p. 12
  - Plant. Aug. 1955, p. 16; Feb. 1956, p. 12; June 1956, p. 12; Feb. 1957, p. 10; Jan. 1958, p. 16
- ## T
- T-shirts. June 1958, p. 5
- Tallow in detergents. Nov. 1956, p. 10
- Tanning leather. June 1957, p. 13; Dec. 1957, p. 4
- Tapeworm. July 1957, p. 10
- Taste test for off-flavors. June 1956, p. 15
- Tax appraisals, improving. Jan. 1957, p. 5
- TBA weedkiller. Feb. 1957, p. 15
- Teaching science. Sept. 1956, p. 2
- Temperature of storage—*see* Frozen foods, Storage, Individual food
- Tenderness:
- Frozen poultry. Sept. 1956, p. 14
  - Meat. Feb. 1957, p. 15
- Tent caterpillar, virus combats. Dec. 1957, p. 7
- Terraces, parallel. June 1956, p. 5
- Tetraploid—*see* Apples, Grapes
- Tetrazolium, versatile research tool. Oct. 1956, p. 5
- Texas Blacklands—*see* Blacklands
- Texas fever tick—*see* Tick, Cattle fever
- Thimet. Feb. 1957, p. 10; Jan. 1958, p. 16
- Thrips, removing from berries. Oct. 1956, p. 15
- Thyroid glands and production. Mar. 1956, p. 13
- Thyroprotein, not for average herd. Nov. 1956, p. 14
- Tick, cattle fever. Dec. 1957, p. 8
- Tillage—*see* Soil
- Timber:
- Large versus small products. Mar. 1958, p. 14
  - Planning log cuts. Sept. 1957, p. 10
  - Present and future resources. Jan. 1956, p. 7
  - Salvaging sawmill wastes. May 1958, p. 14
- Time management. Feb. 1957, p. 16; Mar. 1957, p. 16; May 1957, p. 4
- Time-temperature study of frozen foods. Oct. 1957, p. 12
- Tobacco:
- Dixie Bright variety. Feb. 1956, p. 16
  - Dusting for budworm. Dec. 1956, p. 16
  - Irrigate at top growth period. June 1956, p. 5
  - Wasp allies. July 1956, p. 3
- Tomato:
- Antibiotic for blight. Mar. 1956, p. 10
  - Influences on yield, quality. Jan. 1957, p. 12
  - New varieties. Aug. 1956, p. 12; May 1958, p. 16
  - Powder, other new products. Sept. 1957, p. 11
  - Root-knot nematode. June 1956, p. 12
- Toxicity of pesticides. May 1958, p. 4
- Trace elements—*see* Micronutrients
- Trees:
- Day length. Dec. 1956, p. 10
  - Helpful fungi. Mar. 1956, p. 6; June 1958, p. 3
  - Ornamentals. July 1956, p. 12; May 1957, p. 16
  - Room for fast growth. June 1956, p. 16
  - Saving seedlings. Feb. 1957, p. 7
  - Virus diseases for insect control. April 1956, p. 16; Dec. 1957, p. 7
  - Windbreaks require hardiness. July 1956, p. 7
- Trench silage costs less. Dec. 1957, p. 16
- Tristearin from fats and oils. July 1955, p. 10
- Tristeza disease of citrus. Oct. 1955, p. 8
- Tuberculosis. Aug. 1955, p. 12
- Tulips, bulb storage. Nov. 1957, p. 14
- Tung:
- Nutrition. Aug. 1955, p. 5
  - Oil in paint and varnish. June 1958, p. 6
- Turbinado sugar in candy. Feb. 1956, p. 14
- Turkey:
- Blackhead parasite in cecal worm. Aug. 1957, p. 12
  - Fat in diet. Sept. 1956, p. 13
  - Frozen. Nov. 1955, p. 12



## Turkey—Continued

- Light and male fertility. Oct. 1957, p. 10
- Marketing cooperatives. Mar. 1958, p. 15
- National Improvement Plans. Dec. 1956, p. 15
- Parthenogenesis and fertility. June 1956, p. 3; Nov. 1957, p. 3

Turpentine for synthetic rubber. Mar. 1958, p. 15

## U

Urban survey—see Food

Utilization organization. Apr. 1957, p. 3

## V

- Vacations for farmers. Feb. 1957, p. 16
- Vacuum-cleaner device for seed harvesting. Oct. 1956, p. 16
- Varnish from tung oil. June 1958, p. 6
- Vegetable oil extraction method. May 1956, p. 5
- Vegetative cover for roadsides. Apr. 1957, p. 8
- Vending machines for milk. Jan. 1957, p. 16
- Vertical integration in broiler industry. Mar. 1958, p. 7
- Verticillium wilt:

- Resistance in cotton. Nov. 1955, p. 5
- Saving russet potatoes. July 1955, p. 13

Vesicular exanthema. July 1955, p. 3; Sept. 1955, p. 16

### Virus:

- Against sawflies. Apr. 1956, p. 16
- Controls tent caterpillar. Dec. 1957, p. 7
- Enteritis vaccine for mink. June 1958, p. 12
- Foot-and-mouth. Nov. 1957, p. 15
- In plants. Sept. 1955, p. 15; Oct. 1955, p. 8; Feb. 1956, p. 3; Apr. 1956, p. 7; June 1956, p. 11; Feb. 1958, p. 16;
- See also Barley, Wheat

### Vitamins:

- Amount in plants, influences on. Oct. 1955, p. 3
- B<sub>12</sub> as feed additive. Oct. 1956, p. 10
- B<sub>12</sub> content of dairy products. Oct. 1956, p. 12
- New source of A. Nov. 1957, p. 16
- Utilization of C by human body. Apr. 1957, p. 14

## W

- Warehouse pests of peanuts. Sept. 1957, p. 6
- Wasps, predators for tobacco pests. July 1956, p. 3
- Waste disposal, poultry. Mar. 1957, p. 14
- Wastes, uses for:

- Apple pomace for cattle. May 1958, p. 3
- Pear canning. Apr. 1957, p. 15
- Potatoes, freeze-drying culls. Oct. 1955, p. 10
- Sawmill. May 1958, p. 14
- Sugarcane. July 1956, p. 4; June 1958, p. 16

Water for stock, warming. Feb. 1957, p. 15

### Water management:

- Film as reservoir liner. Dec. 1956, p. 10
- For flood-drought cycles. Nov. 1957, p. 12; Jan. 1958, p. 12
- Moisture, nutrients, and crops. May 1957, p. 6
- Parallel terraces pay. June 1956, p. 5
- Ridge farming. Oct. 1955, p. 7
- Runoff on Great Plains. Apr. 1956, p. 16

## Water management—Continued

- Stored water for sorghum. May 1957, p. 5
- What happens to water. Apr. 1957, p. 6
- See also Irrigation, Soil

Watermelons, improved varieties. Feb. 1958, p. 16

Wax from domestic sugarcane. May 1958, p. 10

### Weather:

- In breeding heat-tolerant cattle. Sept. 1955, p. 11
- Lightning and forest-fire prevention. June 1957, p. 8
- Records warn of potato blight. Jan. 1956, p. 6

### Weeds:

- Can insects fight halogeton? Apr. 1957, p. 11
- Flaming, herbicides for cotton. Apr. 1956, p. 5
- In "clean-cultivated" soybeans. Aug. 1956, p. 16
- Needs in research. Feb. 1957, p. 2
- Shock treatment for weed seeds. Jan. 1957, p. 3
- Weedkillers. Nov. 1955, p. 11; Mar. 1956, p. 5, 10; Apr. 1956, p. 5; Feb. 1957, p. 15; June 1957, p. 15; Apr. 1958, p. 7
- Witchweed. Jan. 1957, p. 15; Apr. 1958, p. 16

Weevils—see Boll weevils

Weigelas, using light to regulate growth. May 1956, p. 16

Weigh-a-day-a-month dairy plan. Sept. 1956, p. 3

Weight grades for dairy cattle. Aug. 1955, p. 13

Weight-height of children. Oct. 1957, p. 13

### Wheat:

- Genetic deviates aid breeding work. May 1956, p. 7
- Hessian-fly control. Dec. 1955, p. 5; Feb. 1956, p. 7
- Leaf-rust resistance. Aug. 1956, p. 8
- Simple tests for stripe mosaic. Feb. 1957, p. 8
- Streak mosaic. July 1957, p. 15
- Stubble mulch beats drought. Feb. 1957, p. 15
- Testing for flour quality. Nov. 1955, p. 15
- Utilization research. Apr. 1956, p. 2

Whipworms in hog pastures. Nov. 1956, p. 15

Windbreaks. Feb. 1956, p. 11; July 1956, p. 7

Wind-erosion control by deep tillage. Sept. 1957, p. 4

Wireworm now resists chlordane. Jan. 1957, p. 13

Witchweed. Jan. 1957, p. 15; Apr. 1958, p. 16

### Wool:

- Breeding for quality, yield. Sept. 15, p. 3
- Color comparator. Apr. 1958, p. 10
- Dye resistance used. May 1957, p. 16
- Fiber structure. Aug. 1956, p. 6
- Improving fabric. Aug. 1956, p. 6
- Shrink resistant. July 1957, p. 6

## Y

Yams, wild, for possible new crop. Jan. 1956, p. 8; Feb. 1958, p. 15

Yeasts in culture collection. Apr. 1958, p. 8

Yeddo hornbeam tree. July 1956, p. 12

## Z

Zebus—see Cattle, heat tolerant

### Zinc:

- For parakeratosis. Apr. 1957, p. 5
- May harm citrus trees. May 1956, p. 10

K38534

